

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
11 July 2002 (11.07.2002)

PCT

(10) International Publication Number
WO 02/053564 A2(51) International Patent Classification⁷: C07D 453/02.
A61K 31/439, A61P 43/00

(21) International Application Number: PCT/EP01/15168

(22) International Filing Date:
20 December 2001 (20.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
P0003130 28 December 2000 (28.12.2000) ES(71) Applicant (for all designated States except US): ALMI-
RALL PRODESFARMA AG [CH/CH]; Lindenhof, Dorf-
strasse 38, CH-6341 Baar (CH).



(72) Inventors; and

(75) Inventors/Applicants (for US only): PRAT QUINONES,

Maria [ES/ES]; Calle Andrea Doria, 2, 1^o 1a, E-08003
Barcelona (ES). FERNANDEZ FORNER, Maria, Do-
lors [ES/ES]; Roger de Flor 221, 5^o, 4a, E-08025 Barcelona
(ES). BUIL ALBERO, Maria, Antonia [ES/ES]; Paris,
50, 1^o, 4a, E-08029 Barcelona (ES).(74) Agents: SRINIVASAN, Ravi, Chandran et al.; J.A.
Kemp & Co., 14 South Square, Gray's Inn, London WC1R
5JJ (GB).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

[Continued on next page]

(54) Title: NOVEL QUINUCLIDINE DERIVATIVES AND MEDICINAL COMPOSITIONS CONTAINING THE SAME



(57) Abstract: A compound of formula (I), wherein B is a phenyl ring, a 5 to 10 membered heteroaromatic group containing one or more heteroatoms, or a naphthalenyl, 5, 6, 7, 8-tetrahydronaphthalenyl, benzo[1,3]dioxolyl, or biphenyl group; R₁, R₂ and R₃ each independently represent a hydrogen or halogen atom, or a hydroxy group, a phenyl group, -OR⁷, -SR⁷, -NR⁷R⁸, -NHCOR⁷, -CONR⁷R⁸, -CN, -NO₂, -COOR⁷ or CF₃ group, or a straight or branched, substituted or unsubstituted lower alkyl group, wherein R⁷ and R⁸ each independently represent a hydrogen atom, a straight or branched lower alkyl group, or together form an

alicyclic ring; or R¹ and R² together form an aromatic or alicyclic ring or a heterocyclic ring. n is an integer from 0 to 4; A represents a group selected from -CH₂-, -CH=CR⁹-, -CR⁹=CH-, -CR⁹R¹⁰-, -CO-, -O-, -S-, -S(O)-, -S(O)₂- and NR⁹, wherein R⁹ and R¹⁰ each independently represent a hydrogen atom, a straight or branched lower alkyl group, or together form an alicyclic ring; m is an integer from 0 to 8, provided that when m=0, A is not -CH₂-; p is an integer from 1 to 2 and the substitution in the azonia bicyclic ring may be in the 2, 3 or 4 position including all possible configurations of the asymmetric carbons; R₄ represents a group of structure: (Formulae II) wherein R¹¹ represents a hydrogen or halogen atom, a hydroxy group, an alkoxy group, a nitro group, a cyano group, -CO₂R¹² or -NR¹²R¹³, wherein R¹² and R¹³ are identical or different and are selected from hydrogen and straight or branched lower alkyl groups, or a straight or branched, substituted or unsubstituted lower alkyl group; R₅ represents an alkyl group of 1 to 7 carbon atoms, an alkenyl group containing 2 to 7 carbon atoms, or a group of formula (III) wherein q=1 or 2 and R¹¹ is as defined above; R₆ represents a hydrogen atom, a hydroxy group, a methyl group or a -CH₂OH group; and X⁻ represents a pharmaceutically acceptable anion of a mono or polyvalent acid.

BEST AVAILABLE COPY

WO 02/053564 A2



European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

- *without international search report and to be republished upon receipt of that report*

NOVEL QUINUCLIDINE DERIVATIVES AND MEDICINAL COMPOSITIONS
CONTAINING THE SAME

This invention relates to new therapeutically useful quinuclidine derivatives, to
5 some processes for their preparation and to pharmaceutical compositions containing
them.

The novel structures according to the invention are antimuscarinic agents with a
potent and long lasting effect. In particular, these compounds show high affinity for
muscarinic M3 receptors. This subtype of muscarinic receptor is present in glands and
10 smooth muscle and mediates the excitatory effects of the parasympathetic system on
glandular secretion and on the contraction of visceral smooth muscle (Chapter 6,
Cholinergic Transmission, in H.P. Rang et al., *Pharmacology*, Churchill Livingstone,
New York, 1995).


M3 antagonists are therefore known to be useful for treating diseases
15 characterised by an increased parasympathetic tone, by excessive glandular secretion
or by smooth muscle contraction (R.M. Eglen and S.S. Hegde, (1997), *Drug News*
Perspect., 10(8):462-469).

Examples of this kind of diseases are respiratory disorders such as chronic
obstructive pulmonary disease (COPD), bronchitis, bronchial hyperreactivity, asthma,
20 cough and rhinitis; urological disorders such as urinary incontinence, pollakiuria,
neurogenic or unstable bladder, cystospasm and chronic cystitis; gastrointestinal
disorders such as irritable bowel syndrome, spastic colitis, diverticulitis and peptic
ulceration; and cardiovascular disorders such as vagally induced sinus bradycardia
(Chapter 7, *Muscarinic Receptor Agonists and Antagonists*, in *Goodman and Gilman's*
25 *The Pharmacological Basis of Therapeutics*, 10th edition, McGraw Hill, New York,
2001).


The compounds of the invention can be used alone or in association with other
drugs commonly regarded as effective in the treatment of these diseases. For example,
they can be administered in combination with β_2 -agonists, steroids, antiallergic drugs,
30 phosphodiesterase IV inhibitors and/or leukotriene D4 (LTD4) antagonists for
simultaneous, separate or sequential use in the treatment of a respiratory disease. The
claimed compounds are useful for the treatment of the respiratory diseases detailed
above in association with β_2 -agonists, steroids, antiallergic drugs or phosphodiesterase
IV inhibitors.

Compounds in related structures have been described as anti-cholinergic and/or anti-spasmodics agents in several patents.

For example, FR 2012964 describes quinuclidinol derivatives of the formula




in which R is H, OH or an alkyl group having 1 to 4 carbon atoms; R¹ is a phenyl or thienyl group; and R² is a cyclohexyl, cyclopentyl or thienyl group, or, when R is H, R¹ and R² together with the carbon atom to which they are attached, form a tricyclic group of the formula:



in which X is -O-, -S- or -CH₂-,
or an acid addition or quaternary ammonium salt thereof.


In United States Patent US 4,465,834 a class of anticholinergic drugs having the formula



are described, in which R¹ is a carbocyclic or branched aliphatic group of 3 to 8 carbon atoms (such as phenyl, cyclohexyl, cyclopentyl, cyclopropyl, cycloheptyl, and isopropyl), R² is a branched or linear aliphatic group containing 3 to 10 carbon atoms


with 1 or 2 olefinic or acetylenic bonds, or is a phenylethynyl, styryl, or an ethynyl group, and R^3 is an alkyl or cyclic group of 4 to 12 carbon atoms containing a tertiary amino nitrogen. The compounds of the invention are also claimed as either the free base or the acid-addition and quaternary ammonium salt forms thereof.

5 In United States Patent 4,843,074 products of formula



are described, wherein $X = H$, halogen, lower alkyl, lower alkoxy, hydroxy and $R =$ morpholinyl, thiomorpholinyl, piperidinyl, 1,4-dioxo-8-azaspiro[4,5]decanyl, 4-(2,6-dimethylmorpholinyl), 4-ketopiperidinyl, 4-hydroxypiperidinyl, 4-substituted piperazinyl. The lower alkyl halide quaternary salts and pharmaceutically acceptable acid addition salts are included in the invention.

US Patent 4,644,003 describes esters of 3-quinuclidinol of alpha disubstituted glycolic acids




and their pharmaceutically acceptable salts,

15 where R is phenyl, unsubstituted or substituted up to three substituents including alkoxy, halogen, nitro, amino, alkylamino, dialkylamino, acylamino, and trifluoromethyl; and wherein R^1 is hydrogen, alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, alkoxyalkyl, cycloalkoxyalkyl, haloalkyl or haloalkenyl.

In WO 92/04346 are described compounds of formula

20




25

and their pharmaceutically acceptable salts, where X is phenyl (optionally substituted) or a thienyl group and "Het" is either (a) a five membered nitrogen-containing heterocyclic group, (b) an oxadiazolyl or thiadiazolyl group, or (c) a six membered nitrogen-containing heterocyclic group, and m is 1 or 2. (For a more detailed description, see the above mentioned publication)

Azoniabicyclic compounds of a general structure related to the compounds of the invention are disclosed in WO 01/04118.

The present invention provides new quinuclidine ester derivatives with potent antagonist activity at muscarinic M3 receptors which have the chemical structure described in formula (I):



wherein B is a phenyl ring, a 5 to 10-membered heteroaromatic group containing one or more heteroatoms, or a naphthalenyl, 5,6,7,8-tetrahydronaphthalenyl, benzo[1,3]dioxolyl, or biphenyl group; R¹, R² and R³ each independently represent a hydrogen or halogen atom, or a hydroxy group, a phenyl group, -OR⁷, -SR⁷, -NR⁷R⁸, -NHCOR⁷, -CONR⁷R⁸, -CN, -NO₂, -COOR⁷ or -CF₃ group, or a straight or branched, substituted or unsubstituted lower alkyl group, wherein R⁷ and R⁸ each independently represent a hydrogen atom, a straight or branched lower alkyl group, or together form an alicyclic ring; or R¹ and R² together form an aromatic or alicyclic ring or a heterocyclic group;


n is an integer from 0 to 4;

A represents a group selected from -CH₂-, -CH=CR⁹-, -CR⁹=CH-, -CR⁹R¹⁰-, -CO-, -O-, -S-, -S(O)-, -S(O)₂- and -NR⁹-, wherein R⁹ and R¹⁰ each independently represent a hydrogen atom, a straight or branched lower alkyl group, or together form an alicyclic ring;


m is an integer from 0 to 8, provided that when m=0, A is not -CH₂-;

p is an integer from 1 to 2 and the substitution in the azonia bicyclic ring may be in the 2,3 or 4 position including all possible configurations of the asymmetric carbons;

R⁴ represents a group of structure:



- 5 wherein R^{11} represents a hydrogen or halogen atom, a hydroxy group, an alkoxy group, a nitro group, a cyano group, $-\text{CO}_2R^{12}$ or $-\text{N}^{R^{12}R^{13}}$, wherein R^{12} and R^{13} are identical or different and are selected from hydrogen and straight or branched lower alkyl groups, or a straight or branched, substituted or unsubstituted lower alkyl group; R^5 represents an alkyl group of 1 to 7 carbon atoms, an alkenyl group containing 2 to 7 carbon atoms, an alkynyl group containing 2 to 7 carbon atoms, a cycloalkyl group of 3 to 7 carbon atoms, or a group of formula



wherein $q = 1$ or 2 and R^{11} is as defined above;

R^6 represents a hydrogen atom, a hydroxy group, a methyl group or a $-\text{CH}_2\text{OH}$ group.

- The asymmetric carbon in the alpha position to the ester group, which is substituted by R^4 , R^5 , and R^6 may have R or S configuration.

X^- represents a pharmaceutically acceptable anion of a mono or polyvalent acid.

- In the quaternary ammonium compounds of the present invention represented by formula (I), an equivalent of an anion (X^-) is associated with the positive charge of the N atom. X^- may be an anion of various mineral acids such as, for example, chloride, bromide, iodide, sulfate, nitrate, phosphate, or an anion of an organic acid such as, for example, acetate, maleate, fumarate, citrate, oxalate, succinate, tartrate, malate, mandelate, trifluoroacetate, methanesulfonate and p-toluenesulfonate. X^- is preferably an anion selected from chloride, bromide, iodide, sulphate, nitrate, acetate, maleate, oxalate, succinate or trifluoroacetate. More preferably X^- is chloride, bromide, trifluoroacetate or methanesulphonate.

- The compounds of the present invention represented by the formula (I) described above, which may have one or more asymmetric carbons, include all the possible stereoisomers. The single isomers and mixtures of the isomers fall within the scope of the present invention.

- As used herein, an alkyl group is typically a lower alkyl group. A lower alkyl group preferably contains 1 to 8, preferably 1 to 6 and more preferably 1 to 4 carbon atoms. In particular it is preferred that such an alkyl group is represented by a methyl, ethyl, propyl, including i-propyl, or butyl including a n-butyl, sec-butyl and tert-butyl group. An alkyl group containing 1 to 7 carbon atoms as mentioned herein, such as

that represented by [REDACTED] may be a C₁₋₄ alkyl group as mentioned above or a straight or branched pentyl, hexyl or heptyl group.

Optionally substituted lower alkyl groups mentioned herein include straight or branched alkyl groups containing from 1 to 6, preferably from 1 to 4, carbon atoms as mentioned above, which may be unsubstituted or substituted in any position by one or more substituents, for example by 1, 2 or 3 substituents. When two or more substituents are present, each substituent may be the same or different. The substituent(s) are typically halogen atoms or hydroxy or alkoxy groups, preferably hydroxy or alkoxy groups.

Alkenyl groups having 2 to 7 carbon atoms mentioned herein, such as those represented by the group R⁵, are straight or branched groups such as ethenyl, or straight or branched propenyl, butenyl, pentenyl, hexenyl or heptenyl. The double bond may be in any position in the alkenyl group, such as on the terminal bond.

Alkynyl groups having 2 to 7 carbon atoms mentioned herein, such as those represented by the group R⁵, are straight or branched groups such as ethynyl, propynyl or straight or branched butynyl, pentynyl, hexynyl or heptynyl. The triple bond may be in any position in the alkynyl group, such as on the terminal bond.

Alkoxy groups mentioned herein, such as those that may be present in the group B, are typically lower alkoxy groups, that is groups containing from 1 to 6 carbon atoms, preferably from 1 to 4 carbon atoms, the hydrocarbon chain being branched or straight. Preferred alkoxy groups include methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, sec-butoxy and t-butoxy.

Cycloalkyl groups and alicyclic groups mentioned herein, unless otherwise specified, typically contain from 3 to 8 carbon atoms, preferably from 3 to 6 carbon atoms. Cycloalkyl groups and alicyclic rings of 3 to 6 carbon atoms include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. Cycloalkyl groups containing from 3 to 7 carbon atoms include cycloalkyl groups of 3 to 6 carbon atoms and cycloheptyl.

The aromatic ring mentioned in relation to R¹ and R² typically contains from 5 to 14, preferably 5 to 10 carbon atoms. Examples of aromatic groups include cyclopentadienyl, phenyl and naphthalenyl.

A heterocyclic or heteroaromatic group mentioned herein is typically a 5 to 10 membered group, such as a 5, 6 or 7 membered group, containing one or more heteroatoms selected from N, S and O. Typically, 1, 2, 3 or 4 heteroatoms are present, preferably 1 or 2 heteroatoms. A heterocyclic or heteroaromatic group may be a single ring or two or more fused rings wherein at least one ring contains a heteroatom. Examples of heterocyclic groups include piperidyl, pyrrolidyl, piperaziny, morpholinyl, thiomorpholinyl, pyrrolyl, imidazolyl, imidazolidinyl, pyrazolinyl, indolinyl,

isoindolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizyl, isoindolyl, indolyl, indazolyl, purinyl, quinolizyl, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalyl, quinazolyl, cinnolyl, pteridinyl, quinuclidinyl, triazolyl, pyrazolyl, tetrazolyl and thienyl. Examples of heteroaromatic groups include pyridyl, thienyl, 5 furyl, pyrrolyl, imidazolyl, benzothiazolyl, pyridinyl, pyrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolyl, indazolyl, purinyl, quinolyl, isoquinolyl, phthalazinyl, naphthyridinyl, quinoxalyl, quinazolyl, cinnolyl, triazolyl and pyrazolyl.

As used herein a halogen atom includes a fluorine, chlorine, bromine or iodine atom, typically a fluorine, chlorine or bromine atom.

10

Preferred compounds of formula (I) are those wherein B represents a phenyl, pyrrolyl, thienyl, furyl, biphenyl, naphthalenyl, 5, 6, 7, 8-tetrahydronaphthalenyl, benzo[1,3]dioxolyl, imidazolyl or benzothiazolyl group, in particular a phenyl, pyrrolyl, thienyl, furyl, biphenyl, naphthalenyl, 5, 6, 7, 8-tetrahydronaphthalenyl or 15 benzo[1,3]dioxolyl group such as a phenyl, thienyl or pyrrolyl group.

B may be unsubstituted or substituted with one, two or three groups (R^1 to R^3) which may be in any position on the ring. Typically it is unsubstituted or substituted with one group, for example when B is a phenyl group it may be substituted in the 2, 3 or 4 position. Typically, R^1 , R^2 and R^3 each independently represent a hydrogen or 20 halogen atom, or a hydroxy, methyl, tert-butyl, $-CH_2OH$, 3-hydroxypropyl, $-OMe$, $-NMe_2$, $-NHCOMe$, $-CONH_2$, $-CN$, $-NO_2$, $-COOMe$ or $-CF_3$ group, in particular a hydrogen, fluorine or chlorine atom or a hydroxy, methyl, $-CH_2OH$, $-OMe$, $-NMe_2$, $-NHCOMe$, $-CONH_2$, $-CN$, $-NO_2$, $-COOMe$ or $-CF_3$ group. The most preferred groups R^1 , R^2 and R^3 are hydrogen, fluorine, chlorine or hydroxy.

25

Examples of substituted phenyl groups which may represent B are tolyl including o-, m- and p-tolyl, 3-cyanophenyl, 2-, 3- and 4-hydroxyphenyl and 2-, 3- and 4-fluorophenyl.

Preferred compounds of formula (I) are those wherein $n = 0$ or 1 ; m is an integer from 1 to 6, particularly 1, 2 or 3; and A represents a $-CH_2-$, $-CH=CH-$, $-CO-$, $-NMe-$, 30 $-O-$ or $-S-$ group, in particular a $-CH_2-$, $-CH=CH-$, $-O-$ or $-S-$ group, for example a $-CH_2-$, $-CH=CH-$ or $-O-$ group.

More preferred salts of formula (I) are those wherein the azoniabicyclo group is substituted on the nitrogen atom with a 3-phenoxypropyl, 2-phenoxyethyl, 3-phenylallyl, phenethyl, 4-phenylbutyl, 3-phenylpropyl, 3-(2-hydroxyphenoxy)propyl, 3-(4- 35 fluorophenoxy)propyl, 2-benzyloxyethyl, 3-pyrrol-1-ylpropyl, 2-thien-2-ylethyl, 3-thien-2-ylpropyl, 3-phenylaminopropyl, 3-(methylphenylamino)propyl, 3-phenylsulfanylpropyl, 3-o-tolylloxypropyl, 3-(2,4,6-trimethylphenoxy)propyl,

3-(2-tert-butyl-6-methylphenoxy)propyl, 3-(4-phenyl-4-yloxy)propyl,
 3-(5,6,7,8-tetrahydronaphthalen-2-yloxy)-propyl, 3-(naphthalen-2-yloxy)propyl,
 3-(naphthalen-1-yloxy)propyl, 3-(2-chlorophenoxy)propyl,
 3-(2,4-difluorophenoxy)propyl, 3-(3-trifluoromethylphenoxy)propyl,
 5 3-(3-cyanophenoxy)propyl, 3-(4-cyanophenoxy)propyl, 3-(3-methoxyphenoxy)propyl,
 3-(4-methoxyphenoxy)propyl, 3-(benzo[1,3]dioxol-5-yloxy)propyl,
 3-(2-carbamoylphenoxy)propyl, 3-(3-dimethylaminophenoxy)propyl,
 3-(4-nitrophenoxy)propyl, 3-(3-nitrophenoxy)propyl, 3-(4-acetylaminoxy)propyl,
 3-(4-methoxycarbonylphenoxy)propyl, 3-[4-(3-hydroxypropyl)phenoxy]propyl,
 10 3-(2-hydroxymethylphenoxy)propyl, 3-(3-hydroxymethylphenoxy)propyl,
 3-(4-hydroxymethylphenoxy)propyl, 3-(2-hydroxyphenoxy)propyl,
 3-(4-hydroxyphenoxy)propyl, 3-(3-hydroxyphenoxy)propyl, 4-oxo-4-thien-2-ylbutyl,
 3-(1-methyl-[1H]-imidazol-2-ylsulfanyl)propyl, 3-(benzothiazol-2-yloxy)propyl,
 3-benzyloxypropyl, 6-(4-phenylbutoxy)hexyl, 4-phenoxybutyl, 4-(4-fluorophenyl)-4-
 15 oxobutyl or 4-oxo-4-phenylbutyl group. Especially preferred salts are those wherein
 the azoniabicyclo group is substituted on the nitrogen atom with a 3-phenoxypropyl,
 2-phenoxyethyl, 3-phenylallyl, phenethyl, 3-phenylpropyl, 3-(3-hydroxyphenoxy)propyl,
 3-(4-fluorophenoxy)propyl, 3-thien-2-ylpropyl group, 4-oxo-4-thien-2-ylbutyl,
 2-benzyloxyethyl, 3-o-tolyloxypropyl, 3-(3-cyanophenoxy)propyl,
 20 3-(methylphenylamino)propyl, 3-phenylsulfanylpropyl, 4-oxo-4-phenylbutyl,
 3-(2-chlorophenoxy)propyl, 3-(2,4-difluorophenoxy)propyl,
 3-(4-methoxyphenoxy)propyl, 3-(benzo[1,3]dioxol-5-yloxy)propyl. Examples of
 especially preferred salts are those wherein the azoniabicyclo group is substituted on
 the nitrogen atom with a 3-phenoxypropyl, 2-phenoxyethyl, 3-phenylallyl, phenethyl, 3-
 25 phenylpropyl, 3-(3-hydroxyphenoxy)propyl, 3-(4-fluorophenoxy)propyl or 3-thien-2-
 ylpropyl group.

Further preferred compounds of formula (I) are those wherein R⁴ represents
 a phenyl, 2-thienyl, 3-thienyl, 2-furyl or 3-furyl group. R¹¹ preferably represents a
 30 hydrogen or halogen atom or an unsubstituted lower alkyl group such as methyl or
 ethyl. Most preferably R¹¹ is a hydrogen atom. Therefore, for example, R⁴ may
 represent an unsubstituted phenyl, 2-thienyl or 2-furyl group. Alternatively, R⁴ may
 represent an unsubstituted phenyl or 2-thienyl group. Preferably, R⁵ represents a
 benzyl, phenethyl, cyclopentyl, cyclohexyl, C₂₋₅ alkyl, C₂₋₅ alkenyl or C₂₋₅ alkynyl group,
 35 particularly a cyclopentyl, cyclohexyl, pentyl, allyl, vinyl, propynyl, phenethyl or benzyl
 group. Alternatively, R⁵ represents a benzyl, cyclopentyl, cyclohexyl, C₂₋₅ alkyl, C₂₋₅
 alkenyl or C₂₋₅ alkynyl group, particularly a cyclopentyl, cyclohexyl, pentyl, allyl, vinyl,

propynyl or benzyl group. The asymmetric carbon alpha to the ester group, which is substituted by R⁴, R⁵ and R⁶ may have R or S configuration.

Preferred compounds of formula (I) are those wherein the group -O-CO-C(R⁴)(R⁵)(R⁶) represents a group selected from 2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl; 2,3-diphenylpropionyloxy; 2-hydroxymethyl-2,3-diphenylpropionyloxy; 2-hydroxy-2,3-diphenylpropionyloxy; 2-hydroxy-3-phenyl-2-thien-2-ylpropionyloxy; 2-hydroxy-2-thien-2-ylpent-4-enoyloxy; 2-hydroxy-2-thien-2-ylheptanoyloxy; 2-hydroxy-2-thien-2-ylpent-3-ynoyloxy; 2-hydroxy-2-thien-2-ylbut-3-enoyloxy; 2-cyclopentyl-2-fur-2-yl-2-hydroxyacetoxyl; 2-cyclohexyl-2-hydroxy-2-phenylacetoxyl; 2-cyclohexyl-2-fur-2-yl-2-hydroxyacetoxyl, 2-cyclopentyl-2-hydroxy-2-phenylacetoxyl, 2-cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl and 2-hydroxy-4-phenyl-2-thien-2-ylbutanoyloxy. Examples of the group -O-CO-C(R⁴)(R⁵)(R⁶) are 2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl; 2,3-diphenylpropionyloxy; 2-hydroxymethyl-2,3-diphenylpropionyloxy; 2-hydroxy-2,3-diphenylpropionyloxy; 2-hydroxy-3-phenyl-2-thien-2-ylpropionyloxy; 2-hydroxy-2-thien-2-ylpent-4-enoyloxy; 2-hydroxy-2-thien-2-ylheptanoyloxy; 2-hydroxy-2-thien-2-ylpent-3-ynoyloxy; 2-hydroxy-2-thien-2-ylbut-3-enoyloxy; 2-cyclopentyl-2-fur-2-yl-2-hydroxyacetoxyl; 2-cyclohexyl-2-hydroxy-2-phenylacetoxyl and 2-cyclohexyl-2-fur-2-yl-2-hydroxyacetoxyl.

In the most preferred embodiments of the invention p is 2, the azoniabicyclo[2.2.2]octane ring is substituted in the 3 position, and this substituted carbon atom has (R)-configuration.

The following compounds are intended to illustrate but not to limit the scope of the present invention:

- (3R)-3-(2,3-Diphenylpropionyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2,3-Diphenylpropionyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2*)-2-Hydroxymethyl-2,3-diphenylpropionyloxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 1)
- (3R)-3-[(2*)-2-Hydroxymethyl-2,3-diphenylpropionyloxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 1)
- (3R)-3-[(2*)-2-Hydroxymethyl-2,3-diphenylpropionyloxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 2)

- (3R)-3-[(2*)-2-Hydroxy-2-methyl-2,3-diphenylpropionyloxy]-1-phenyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 2)
- (3R)-3-[(2*)-2-Hydroxy-2,3-diphenylpropionyloxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide (diastereomer 1)
- 5 (3R)-3-[(2*)-2-Hydroxy-2,3-diphenylpropionyloxy]-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide (diastereomer 1)
- (3R)-3-[(2*)-2-Hydroxy-2,3-diphenylpropionyloxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 2)
- (3R)-3-[(2*)-2-Hydroxy-2,3-diphenylpropionyloxy]-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 2)
- 10 (3R)-3-(2-Hydroxy-3-phenyl-2-thien-2-ylpropionyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-3-phenyl-2-thien-2-ylpropionyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 (3R)-3-(2-Hydroxy-2-thien-2-ylpent-4-enoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-(2-Hydroxy-2-thien-2-ylpent-4-enoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-2-thien-2-ylpent-4-enoyloxy)-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide
- 20 (3R)-3-(2-Hydroxy-2-thien-2-ylheptanoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-2-thien-2-ylheptanoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 (3R)-3-(2-Hydroxy-2-thien-2-ylpent-3-ynoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-2-thien-2-ylpent-3-ynoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-2-thien-2-ylbut-3-enoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 (3R)-3-(2-Hydroxy-2-thien-2-ylbut-3-enoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- 35 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(2-benzyloxyethyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenylallyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(4-fluorophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 10 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(4-oxo-4-thien-2-ylbutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[4-(4-fluorophenyl)-4-oxobutyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(3-hydroxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 1-(2-Benzyloxyethyl)-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-o-tolyloxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(3-Cyanophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(methylphenylamino)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(4-oxo-4-phenylbutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(2,4,6-trimethylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 1-[3-(2-Chlorophenoxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 35 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(3-trifluoromethylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

- 1-[3-(Biphenyl-4-yloxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(2,4-difluorophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-methoxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(5,6,7,8-tetrahydronaphthalen-2-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 10 1-[3-(Benzo[1,3]dioxol-5-yloxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 1-[3-(2-Carbamoylphenoxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(3-dimethylaminophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 1-[3-(4-Acetylaminophenoxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-methoxycarbonylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-nitrophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-hydroxymethylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- 25 (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenylallyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 35 (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-fluorophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

- (3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 10 (3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 (3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 (3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 (3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 (3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 35 (3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

- (3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxyl-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxyl-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxyl-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxyl-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxyl-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 10 (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxyl-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxyl-1-(3-phenylsulfanypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxyl-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxyl-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxyl-1-(3-phenylsulfanypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 (3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 1-[3-(3-Cyanophenoxy)propyl]-(3R)-3-[(2S)-2-cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 1-[3-(3-Cyanophenoxy)propyl]-(3R)-3-[(2R)-2-cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-4-phenyl-2-thien-2-ylbutanoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 35 (3S)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide

(3S)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate


(3S)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

5 4-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

4-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

10 ((*): Configuration not assigned; either the (2R)- or the (2S)- isomers of the above compounds may be produced).

The present invention also provides processes for preparing compounds of formula (I). The quaternary ammonium derivatives of general formula (I), may be prepared by reaction of an alkylating agent of general formula (II) with compounds of general formula (III). In formulae (I), (II) and (III), R^1 , R^2 , R^3 , B, A, X^- , R^4 , R^5 , R^6 , n, m and p are as defined above.



20

In formula (II), W represents any suitable leaving group, such as a group X as defined above. Preferably, W represents a group X.


This alkylation reaction may be carried out by two different experimental procedures, a) and b) which are described above. In particular method b) provides a new experimental process, using solid phase extraction methodologies, that allows the parallel preparation of several compounds. Methods a) and b) are described in the

25

experimental section. W represents a group other than X, the quaternary ammonium salt of formula (I) is produced from the product of method (a) or (b) by carrying out an exchange reaction according to standard methods to replace the anion W^- with the desired anion X^- .

5

Compounds of general formula (II) which are not commercially available have been prepared by synthesis according to standard methods. For example, compounds wherein $n = 0$ and $A = -O-$, $-S-$ or $-NR^9-$, wherein R^9 is as defined above, were obtained by reaction of the corresponding aromatic derivative or its potassium salt with an alkylating agent of general formula $Y-(CH_2)_m-X$, wherein X may be a halogen and Y may be a halogen or a sulphonate ester. In other examples, compounds of general formula (II), where $n \geq 1$ were synthesised from the corresponding alcohol derivative of general formula (IV) by known methods



15


wherein R^1 , R^2 , R^3 , n , m , and A are as defined above.

Compounds of general formula (III) may be prepared by different methods. These procedures are illustrated in the following schemes and detailed in the experimental section.

Method c)

20

Compounds of general formula (III) may be synthesised by transesterification of a compound of formula (V) with a compound of formula (VI).



25


in which formulae R^4 , R^5 , R^6 and p are as defined above and L represents a leaving group. For example L may be a chlorine atom, an imidazol-1-yl group, or a group –

OR¹⁴ wherein R¹⁴ represents a straight, or branched, substituted or unsubstituted lower alkyl group or a -COR¹⁵ group wherein R¹⁵ represents -COCR¹⁶R¹⁷. Typically L is -OR¹⁴ wherein R¹⁴ is methyl, ethyl or propyl, or L is an imidazol-1-yl group.

Intermediates of formula (VI) may be prepared by standard methods described in the literature, for example as in FR 2012964.

Method d)


Compounds of formula (III) where R⁶ is a hydroxy group and p, R⁴ and R⁵ are as described above, may also be prepared from the glyoxalate esters of general formula (VII) by reaction with the corresponding Grignard reagent.



Compounds of general formula (VII) may be prepared by standard methods described in the literature (WO 01/04118; WO 92/04346)


Method e)

Compounds of formula (III) where R⁶ is a group -CH₂OH, and p, R⁴ and R⁵ are as described above, may also be prepared from the corresponding compound of formula (III), where R⁶ is an hydrogen atom, by reaction with formaldehyde in basic conditions. (Organic Syntheses Collective Volumes, VII, 271-274, (1990); WO 93/06098)



As will be described in the experimental part, the diastereomers of the compounds of formula (III) may be separated by conventional methods, for example by column chromatography or crystallisation.

The following compounds are compounds of general formula (III) which have not been described before:
compounds of formula (III)



(III)

wherein p, R⁴, R⁵ and R⁶ are as defined above, the substituent on the azabicyclo group is at position 3 or 4 and when it is at position 3 this substituted carbon has an enantiomerically pure R or S configuration, provided that when R⁴ is a 3-thienyl group and R⁵ is a cyclohexyl group R⁶ is not a hydroxy group.

The substituent on the azabicyclo group is preferably at position 3 and may have R or S configuration. The carbon substituted by R⁴, R⁵ and R⁶ may have R or S configuration. The compound may be a single isomer.

Examples of the new compounds of formula (III) include:

(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester ($[\alpha]^{22}_D = +19.7^\circ$ (c=1, CHCl₃)).

(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester ($[\alpha]^{22}_D = -14.2^\circ$ (c=1, CHCl₃)).

(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester ($[\alpha]^{22}_D = +21.1^\circ$ (c=1, CHCl₃)).

(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester ($[\alpha]^{22}_D = -23.5^\circ$ (c=1, CHCl₃)).

(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid 1-azabicyclo[2.2.2]oct-4-yl ester ($[\alpha]^{22}_D = -27.6^\circ$ (c=1, CHCl₃)).

In particular:

(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester ($[\alpha]^{22}_D = +19.7^\circ$ (c=1, CHCl₃)).

(2R)-2-Cyclopentyl-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester ($[\alpha]^{22}_D = -14.2^\circ$ (c=1, CHCl₃)).

(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester ($[\alpha]^{22}_D = +21.1^\circ$ (c=1, CHCl₃)).

- 5 (2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester ($[\alpha]^{22}_D = -23.5^\circ$ (c=1, CHCl₃)).

Compounds of formula (V) could be:

- 10 4-hydroxy-1-azabicyclo[2.2.1]heptane, described in WO93/15080
 4-hydroxy-1-azabicyclo[2.2.2]octane, described in Grob, C.A. et.al. *Helv.Chim.Acta* (1958), 41, 1184-1190
 (3R)-3-hydroxy-1-azabicyclo[2.2.2]octane or (3S)-3-hydroxy-1-azabicyclo[2.2.2]octane, described in Ringdahl, R. *Acta Pharm Suec.* (1979), 16, 281-283 and commercially
 15 available from CU Chemie Uetikon GmbH.

The following examples are intended to illustrate, but not to limit, the experimental procedures that have been described above.

- The structures of the prepared compounds were confirmed by ¹H-NMR and MS.
 20 The NMR spectra were recorded using a Varian 300 MHz instrument and chemical shifts are expressed as parts per million (δ) from the internal reference tetramethylsilane. Their purity was determined by HPLC, using reverse phase chromatography on a Waters instrument, with values greater than 95% being obtained. Molecular ions were obtained by electrospray ionization mass spectrometry on a
 25 Hewlett Packard instrument. Optical rotations were obtained using a PERKIN-ELMER 241 MC Polarimeter.

Method -a-

Example 13

- 30 **Preparation of (3R)-3-(2-Hydroxy-2-thien-2-ylpent-4-enoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide**
 250 mg (0.81 mmol) of 2-Hydroxy-2-thien-2-ylpent-4-enoic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester, Intermediate I-3, were dissolved in 5 ml of acetonitrile and 7.5 ml of chloroform. To this solution were added 0.63 ml (4.1 mmol) of
 35 phenoxypropylbromide. After stirring for 48 h at room temperature under a N₂ atmosphere, solvents were evaporated. Ether was added and the mixture stirred. The

solid obtained was dried and washed several times with ether. The yield was 0.3 g (71%) of the title compound as a mixture of diastereomers; mp: 137° C.

¹H-NMR (DMSO-d₆): δ 1.70-2.05 (m, 4H), 2.05-2.35 (m, 3H), 2.70-2.83 (m, 1H), 2.90-3.02 (m, 1H), 3.25-3.60 (m, 7H), 3.82-3.97 (m, 1H); 3.97-4.10 (m, 2H), 5.05-5.25 (m, 3H), 5.70-5.90 (m, 1H), 6.50 (d, 1H, OH), 6.90-7.05 (m, 4H), 7.10-7.20 (m, 1H), 7.27-7.35 (m, 2H), 7.45 (m, 1H).

MS : [M-Br]⁺: 442.

Method - b -

10 Example 35

Preparation of (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

20 mg (0.06 mmols) of (2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester (Intermediate I-15a), were dissolved in 1ml of DMSO. To this solution 66 mg (0.30 mmol) of 3-(naphthalen-1-yloxy)propyl chloride were added. After stirring overnight at room temperature, the mixture was purified by solid phase extraction with a cation exchange Mega Bond Elut cartridge, previously conditioned at pH = 7.5 with 0.1 M NaH₂PO₄ buffer. The reaction mixture was applied to the cartridge and washed first with 2 ml of DMSO and then three times with 5 ml of CH₃CN, rinsing away all starting materials. The ammonium derivative was eluted with 5 ml of 0.03 M TFA solution in CH₃CN:CHCl₃ (2:1). This solution was neutralized with 300 mg of poly(4-vinylpyridine), filtered and evaporated to dryness. The yield was 10 mg (26 %) of title compound.

¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m, 8H), 1.80-2.10 (m, 4H), 2.20-2.37 (m, 3H), 2.75-2.92 (m, 1H), 3.20-3.65 (m, 7H), 3.90-4.05 (m, 1H), 4.15-4.30 (m, 2H), 5.15-5.22 (m, 1H), 6.24 (s, 1H, OH), 6.95-7.05 (m, 2H), 7.15-7.20 (m, 1H), 7.40-7.60 (m, 5H), 7.85-7.95 (m, 1H), 8.20-8.25 (m, 1H).

MS: [M-CF₃COO]⁺ : 520

30 The spatial configurations of the compounds of general formula (III) have been deduced from the configurations of their corresponding acids. These were determined either comparing the values of [α] obtained with the values described in the literature or applying the Circular Dichroism (CD) technique.

Since the CD curve of (2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid is known (A. Tambuté and A. Collet, Bulletin de la Société Chimique de France, 1984, N° 1-2, pages II77 to II82) and all the acids evaluated are structurally very similar to (2S)-

2-Cyclohexyl-2-hydroxy-2-phenylacetic acid, it can be assumed that the relative positions of the aryl, cycloalkyl, carboxyl and hydroxyl substituents may be identified by comparison of their respective $\Delta\epsilon$ values. The resulting configurations have been expressed as R or S according to Cahn-Ingol-Prelog system. (see TABLE 1)

- 5 The curves of CD spectra were recorded with a Jasco-720 spectrophotometer (Software J-700) from 0.43 mM MeOH solutions of the samples in 1 mm cells at 25°C.

TABLE 1

Compound	Circular Dichroism			Configuration
	$[\alpha]^{22}_D$	λ (nm)	$\Delta\epsilon$ ($M^{-1}cm^{-1}$)	
(+)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid	+ 23.1° (c=1.4, EtOH)	224	+12.1	S ^a
(-)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid	- 23.6° (c=1.4, EtOH)	224	-11.7	R
(-)-2-Cyclopentyl-2-hydroxy-2-phenylacetic acid	-1.52° (c=3, MeOH)	224	-8.93	R ^b
(+)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid ^{c,1}	+ 6.63° (c=1, EtOH)	233	+4.18	R
(-)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid ^{c,2}	-6.44° (c=1, EtOH)	233	-4.19	S
(-)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetic acid ^{c,3}	-15.1° (c=1, EtOH)	235	-5.40	S
(+)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid	+31.95° (c=1, EtOH)	230	+7.64	S
(-)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid	-32.10° (c=1, EtOH)	230	-7.44	R
(+)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid	+ 40.9° (c=1, EtOH)	230	+10.9	S
(-)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid	-39.7° (c=1, EtOH)	230	-10.8	R
^a A. Tambuté et A. Collet; Bulletin de la Société Chimique de France, (1984), N°1-2, II-77-II-82: Configuration S assigned: $[\alpha]^{24}_D = + 25.2^\circ$ (c=1.4, EtOH), $\Delta\epsilon = +12.9 M^{-1}cm^{-1}$ ($\lambda = 225$ nm)				
^b M. Mitsuya et al.; Bioorg. Med. Chem., (1999), Vol 7, 2555-2567 : Configuration R assigned $[\alpha]^{20}_D = -1.9^\circ$ (c=3, MeOH)				
^c E. Atkinson et al. J. Med. Chem., (1977), Vol 20, N° 12, 1612-1617. The values for $[\alpha]$ were given (Configuration not assigned); c,1 and -c,2 : + 51.3° and - 51.0° (rotations observed at 350 nm (c= 2-5%, MeOH));				

Method - c -

Methyl ester derivatives of general formula (VI) were prepared by standard methods described in the literature or following the procedures described in the examples:

5 Intermediates I-9, I-10, I-11, I-12, I-13.

Intermediate I-1**Preparation of 2-Hydroxy-2-thien-2-ylheptanoic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester**

10 2.7 g of 2-Hydroxy-2-thien-2-ylheptanoic acid methyl ester (Intermediate I-9) (0.011mol) were dissolved in 70 ml of toluene. To this solution were added 1.63 g (0.0128 mol) of (3R)-3-hydroxy-1-azabicyclo[2.2.2]octane and 0.18 g (0.0045 mol) of HNa (60% dispersion in mineral oil). The mixture was refluxed with continuous removal of distillate with replacement with fresh toluene when necessary for 1 hour. The cooled
15 mixture was extracted with 1N HCl acid, the aqueous layer washed with ether, basified with K₂CO₃ and extracted with CHCl₃. The organic layer was washed with water, dried over MgSO₄ and evaporated. The yield was 2.85 g (76%) of the title product as a mixture of diastereomers, structure confirmed by ¹H-NMR.

¹H-NMR (CDCl₃): δ 0.80-0.95 (m, 3H), 1.20-1.40 (m, 6H), 1.40-1.90 (m, 4H), 1.95- 2.25
20 (m, 3 H), 2.50 - 2.95 (m, 5H), 3.10-3.30 (m, 1H), 4.4 (bs, 1H, OH), 4.82-4.94 (m, 1H), 6.94-7.02 (m, 1H), 7.06-7.14 (m, 1H), 7.20-7.26 (m, 1H).

MS: [M+1]⁺: 338.

(Compound also prepared following method d)

25

Intermediate I-2**Preparation of 2,3-Diphenylpropionic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester**

Prepared using the same method as for Intermediate I-1, but from 2,3-diphenylpropionic acid methyl ester. The yield was 1.71 g (61.5%) of the title product
30 as a mixture of diastereomers.

¹H-NMR (DMSO): δ 1.05-1.20 (m, 1H), 1.30-1.60 (m, 3H), 1.65-1.75 (m, 1H), 2.10-2.20
(m, 1H), 2.30-2.70 (m, 4H), 2.85-3.10 (m, 2H), 3.20-3.40 (m, 1H), 3.95-4.10 (m, 1H),
4.50-4.65 (m, 1H), 7.10-7.45 (m, 10H).

MS: [M+1]⁺: 336

35 (2,3-diphenylpropionic acid methyl ester may be prepared from the 2,3-diphenylpropionic acid, commercially available)

Intermediate I-3

Preparation of 2-Hydroxy-2-thien-2-ylpent-4-enoic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

Prepared using the same method as for Intermediate I-1, but from 2-Hydroxy-2-thien-2-ylpent-4-enoic acid methyl ester (Intermediate I-10). The yield was 1.76 g (63.1%) of the title product as a mixture of diastereomers.

¹H-NMR (CDCl₃): δ 1.25-1.80 (m, 4H), 1.80-2.10 (m, 1H), 2.50-3.05 (m, 7H), 3.10-3.35 (m, 2H), 4.55 (bs, 1H, OH), 4.75-4.95 (m, 1H), 5.10-5.30 (m, 2H), 5.70-5.95 (m, 1H), 6.95-7.05 (m, 1H), 7.10-7.20 (m, 1H), 7.20-7.30 (m, 1H).

MS: [M+1]⁺: 308

(Compound also prepared following method d)

Intermediate I-4

Preparation of 2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

Prepared using the same method as for Intermediate I-1, but from 13.73 g (0.057 mol) of 2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid methyl ester (Intermediate I-11) dissolved in 350 ml of toluene, 8.6 g (0.067 mol) of (3R)-3-hydroxy-1-azabicyclo[2.2.2]octane and 1.37 g (0.0342 mol) of HNa (60% dispersion in mineral oil). The oil obtained (10.33 g) was purified by chromatography on silica gel eluting with chloroform/methanol/ammonia 97:3:0.3. Appropriate fractions were combined and evaporated to obtain the two diastereomers: I-4a and I-4b.

Intermediate I-4a

(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester

1.59 g of an oil corresponding to the first eluted diastereomer was triturated with a mixture of ethyl ether / isopropyl ether 1:1, to give 0.82 g (8.6 %, based on single isomer) of a solid whose structure was confirmed by ¹H-NMR as a pure diastereomer.

[α]_D²² = + 21.1° (c=1, CHCl₃).

¹H-NMR (CDCl₃): δ 1.10-1.45 (m, 8H), 1.45-1.60 (m, 2H), 1.60-1.85 (m, 4H), 1.94-2.02 (m, 1H), 2.26-2.38 (m, 1H), 2.70-2.92 (m, 5H), 3.20-3.28 (m, 1H), 3.78 (bs, 1H, OH), 4.90 (m, 1H), 6.30-6.40 (m, 2H), 7.40 (m, 1H).

MS: [M+1]⁺: 334.

Intermediate I-4a was hydrolysed (EtOH/NaOH 2N, 2h r.t, 1h 60°C), to yield (+)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid : $[\alpha]_D^{22} = +40.9^\circ$ (c=1, EtOH). Configuration S was assigned (See TABLE 1)

Intermediate I-4b

5 (2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester

1.12 g of an oil corresponding to the second eluted diastereomer was triturated with a mixture of ethyl ether / isopropyl ether 1:1, to give 0.57 g (6 %, based on single isomer) of a solid whose structure was confirmed by $^1\text{H-NMR}$ as a pure diastereomer.

10 $[\alpha]_D^{22} = -23.5^\circ$ (c=1, CHCl_3).

$^1\text{H-NMR}$ (CDCl_3): δ 1.10-1.50 (m, 8H), 1.50-1.90 (m, 6H), 2.04-2.12 (m, 1H), 2.24-2.36 (m, 1H), 2.46-2.58 (m, 1H), 2.68-2.94 (m, 4H), 3.12-3.22 (m, 1H), 3.77 (bs, 1H, OH), 4.90 (m, 1H), 6.40 (m, 2H), 7.42 (m, 1H)

MS: $[\text{M}+1]^+$: 334.

15 Intermediate I-4b was hydrolysed (EtOH/NaOH 2N, 2h r.t, 1h 60°C), to yield (-)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid : $[\alpha]_D^{22} = -39.7^\circ$ (c=1, EtOH). Configuration R was assigned. (See TABLE 1)

Intermediate I-5

20 Preparation of 2-Cyclopentyl-2-hydroxy-2-phenylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester

Prepared using the same method as for Intermediate I-1, but from 13.5 g (0.0576 mol) of 2-Cyclopentyl-2-hydroxy-2-phenylacetic acid methyl ester (commercially available) dissolved in 350 ml of toluene, 8.0 g (0.063 mol) of (3R)-3-hydroxy-1-azabicyclo[2.2.2]octane and 0.91 g (0.023 mol) of HNa (60% dispersion in mineral oil).
25 The yield was 13.1 g (69%) of the title product as an oil mixture of diastereomers, structure confirmed by $^1\text{H-NMR}$. The two diastereomers I-5a and I-5b were separated after several crystallizations.

Intermediate I-5a

30 (2R)-2-Cyclopentyl-2-hydroxy-2-phenylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

Several treatments of the oil mixture of diastereomers with different mixtures of diethyl ether/hexane and diisopropyl ether/hexane (cooling at -60°C) yield 4.3 g of a white solid identified by $^1\text{H-NMR}$ as an enriched diastereomer I-5a. This solid was
35 recrystallised twice from diethyl ether/hexane (cooling at 0°C) to yield 2 g (21%) of pure diastereomer.

$$[\alpha]^{22}_D = -10.8^\circ (c=1, \text{CHCl}_3)$$

¹H-NMR (CDCl₃): δ 1.25-1.80 (m, 11H), 1.80-1.95 (m, 1H), 2.05-2.15 (m, 1H), 2.40-2.50 (m, 1H), 2.62-3.05 (m, 5H), 3.05-3.18 (m, 1H), 3.80 (s, 1H, OH), 4.85-4.90 (m, 1H), 7.22-7.42 (m, 3H), 7.60-7.75 (m, 2H).

- 5 Intermediate I-5a was hydrolysed (EtOH/NaOH 2N, 2h r.t. and 2h at 60°C) to give (-)-2-Cyclopentyl-2-hydroxy-2-phenylacetic acid as a pure enantiomer: $[\alpha]^{22}_D = -1.52^\circ (c=3, \text{MeOH})$. This value was assigned to the R configuration provided that in the literature (M. Mitsuya et al.; Bioorg. Med. Chem., (1999), 7, 2555-2567) the R enantiomer has been described with $[\alpha]^{20}_D = -1.9^\circ (c=3, \text{MeOH})$. (See TABLE 1)

10 **Intermediate I-5b**

(2S)-2-Cyclopentyl-2-hydroxy-2-phenylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

- The mother liquors from the first solid I-5a were evaporated and treated with maleic acid in isopropanol/diethyl ether. After cooling at 0-5°C, 7.0 g of a white solid were
15 obtained and identified by ¹H-RMN as the maleate salt of a mixture enriched with the second diastereomer I-5b. After three crystallisations of this product from acetonitrile/diethyl ether (1:2.2), 2.4 g (18.7%, based on free base) of the maleate salt enriched with the second diastereomer I-5b (in a proportion 88:12 as determined by H-RMN) were obtained.

- 20 This maleate salt enriched with the second diastereomer I-5b (88:12) was treated with CHCl₃ and K₂CO₃ solution to obtain the free base.

I-5b (free base):

$$[\alpha]^{22}_D = +19.5^\circ (c=1, \text{CHCl}_3)$$

- ¹H-NMR (CDCl₃): δ 1.20-1.80 (m, 12H), 1.85-2.0 (m, 1H), 2.60-3.05 (m, 6H), 3.20-3.35
25 (m, 1H), 3.80 (s, 1H, OH), 4.75-4.82 (m, 1H), 7.20-7.45 (m, 3H), 7.55-7.75 (m, 2H).

The signals corresponding to the diastereomer I-5a (12%) were observed at 2.05-2.15, 2.40-2.50, 3.05-3.18, 4.85-4.90 ppm.

Configuration S was assigned in view of the results obtained for Intermediate I-5a. (See TABLE 1).

30

Intermediate I-6

Preparation of 2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

- Prepared using the same method as for Intermediate I-1, but from 16.2 g (0.064 mol) of
35 2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetic acid methyl ester (Intermediate I-12), dissolved in 400 ml of toluene, 9.5 g (0.074 mol) of (3R)-3-hydroxy-1-

azabicyclo[2.2.2]octane and 1.51 g (0.038 mol) of HNa (60% dispersion in mineral oil). The oil obtained (10.97 g) was purified by chromatography on silica gel, eluting with chloroform/methanol/ammonia 95:5:0.5, to obtain 8.97 g of a pure product as a mixture of diastereomers, structure confirmed by ¹H-RMN. Three crystallisations of this mixture from ethyl ether yield 1.68 g (15.2%) of a pure diastereomer (Intermediate I-6a). The mother liquors of the crystallisations were enriched with the other diastereomer (Intermediate I-6b).

Intermediate I-6a

(2S)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

$[\alpha]^{22}_D = -16.5^\circ$ (c=1, CHCl₃).

¹H-NMR (CDCl₃): δ 1.05-1.85 (m, 13H), 1.85-2.0 (m, 1H), 2.0-2.16 (m, 2H), 2.58-2.68 (m, 1H), 2.70-3.0 (m, 4H), 3.14-3.24 (m, 1H), 4.0 (s, 1H, OH), 4.90-5.0 (m, 1H), 6.95-7.05 (m, 1H), 7.10-7.15 (m, 1H), 7.20-7.30 (m, 1H).

MS: [M+1]⁺: 350.

Intermediate I-6a was hydrolysed (EtOH/NaOH 2N, 2h r.t, 1h 60°C), to yield (-)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetic acid : $[\alpha]^{22}_D = -15.1^\circ$ (c=1, EtOH).

Configuration S was assigned. (See TABLE 1).

Intermediate I-6b

(2R)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3(R)-yl ester.

¹H-NMR (CDCl₃): δ 1.05-2.0 (m, 14H), 2.0-2.20 (m, 2H), 2.65-3.02 (m, 5H), 3.24-3.36 (m, 1H), 4.0 (s, 1H, OH), 4.80-4.90 (m, 1H), 6.95-7.05 (m, 1H), 7.10-7.20 (m, 1H), 7.20-7.30 (m, 1H).

The signals corresponding to the Intermediate I-6a (aprox. 25%) were observed at 2.58-2.68, 3.14-3.24 and 4.90-5.0 ppm.

MS: [M+1]⁺: 350.

Configuration R was assigned in view of the results obtained with Intermediate I-6a. (See TABLE 1).

Intermediate I-7

(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid (3S)-1-azabicyclo[2.2.2]oct-3-yl ester.

510 mg (0.00225 mol) of (2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid (obtained by hydrolysis of Intermediate I-15a) were dissolved in 7 ml of DMF. This solution was stirred at room temperature and 638 mg (0.00393 mol) of

1,1'-carbonyldiimidazole were added in several portions. After 4 h the reaction mixture was cooled to 0°C and 315 mg (0.00248 mol) of (3S)-3-hydroxy-1-azabicyclo[2.2.2]octane and 83 mg (0.0021 mol) of HNa (60% dispersion in mineral oil) were added thereto. After stirring 112 h at room temperature the reaction mixture was treated with water and extracted three times with diethyl ether. The organic layers were combined, washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated and the residue was purified by silica gel column chromatography (eluent CHCl₃/MeOH 15:1) to obtain 360 mg (47.6%) of the title product as an oil, structure confirmed by ¹H-RMN.

10 $[\alpha]_D^{22} = -18.16^\circ$ (c=1, CHCl₃)

¹H-NMR (CDCl₃): δ (Same description as in Intermediate I-15b)

MS: [M+1]⁺: 336.

Intermediate I-8

15 **(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid 1-azabicyclo[2.2.2]oct-4-yl ester.**

Prepared using the same method as for Intermediate I-7, but from a solution of 660 mg (0.00282 mol) of (2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid (obtained by hydrolysis of Intermediate I-16a) in 9 ml of DMF, 548 mg (0.00338 mol) of 1,1'-carbonyldiimidazole, 394 mg (0.0031 mol) of 4-hydroxy-1-azabicyclo[2.2.2]octane and 104 mg (0.00259 mol) of HNa (60% dispersion in mineral oil). After 44 h of stirring at room temperature the reaction mixture was treated with water and extracted three times with diethyl ether. The organic layers were combined, washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated and the residue was purified by silica gel column chromatography eluting with CHCl₃ to CHCl₃/MeOH 15:1. The yield was 300 mg (31 %) of the title product.

25 $[\alpha]_D^{22} = -27.6^\circ$ (c=1, CHCl₃)

¹H-NMR (CDCl₃): δ 1.0-1.55 (m, 7H), 1.55-1.75 (m, 2H), 1.75-1.85 (m, 1H), 1.85-2.05 (m, 6H), 2.10-2.22 (m, 1H), 2.90-3.10 (m, 6H), 3.60-3.80 (bs, 1H, OH), 7.20-7.40 (m, 3H), 7.57-7.67 (m, 2H).

30 MS: [M+1]⁺: 344.

Intermediate I-9

Preparation of 2-Hydroxy-2-thien-2-ylheptanoic acid methyl ester.

35 50 ml of a ether solution of 0.0338 mol of pentylmagnesium bromide prepared from 5.1 g of 1-bromopentane (0.0338 mol) and 0.0372 mols of magnesium, were added to

a solution of 5 g of 2-oxo-2-thien-2-ylacetic acid methyl ester dissolved in 40 ml of a solution of ether/THF (50:50), at -70°C under a N_2 atmosphere. The mixture was stirred at this temperature for 10 minutes, and then warmed to room temperature. After 16 h, the reaction mixture was treated with a saturated solution of ammonium chloride and extracted three times with ethyl acetate. The organic phases were combined, washed with water, and dried over MgSO_4 . After removal of the solvent, the oil obtained was purified by column chromatography (silica gel) using mixtures of hexane/AcOEt (25:1 to 15:1) as eluent. The yield was 2.7 g (38%) of a pure product, whose structure was confirmed by $^1\text{H-NMR}$.

$^1\text{H-NMR}$ (CDCl_3): δ 0.80-1.0 (m, 3H), 1.10-1.45 (m, 6H), 1.90-2.30 (m, 2H), 3.80 (s, 3H), 4.05 (s, 1H, OH), 7.0 (m, 1H), 7.10 (m, 1H), 7.30 (m, 1H).

(2-Oxo-2-thien-2-ylacetic acid methyl ester was prepared from the commercially available 2-oxo-2-thien-2-ylacetic acid by a standard method).

Intermediate I-10

Preparation of 2-Hydroxy-2-thien-2-ylpent-4-enoic acid methyl ester

Prepared using the same method as for Intermediate I-9. The yield was 1.92 g. 45.3 %.

$^1\text{H-NMR}$ (CDCl_3): δ 2.75-3.0 (m, 2H), 3.80 (s, 3H), 4.0 (s, 1H, OH), 5.10-5.30 (m, 2H), 5.70-5.90 (m, 1H), 6.95-7.05 (m, 1H), 7.10-7.20 (m, 1H), 7.25-7.35 (m, 1H).

MS: $[\text{M}]^+ = 212$.

Intermediate I-11

Preparation of 2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid methyl ester.

Prepared using the same method as for Intermediate I-9, but from 73 ml (0.146 mol) of a solution of cyclohexylmagnesium chloride 2M in diethyl ether and 22.51 g (0.146 mol) of 2-fur-2-yl-2-oxoacetic acid methyl ester (dissolved in 330 ml of THF). The oil obtained was purified by column chromatography (silica gel) using a mixture of hexane/AcOEt 9:1 as eluent. After removal of the solvent 13.73 g (39%) of a pure product were obtained whose structure was confirmed by MS and $^1\text{H-NMR}$.

$^1\text{H-NMR}$ (CDCl_3): δ 1.05-1.75 (m, 6H), 1.75-1.95 (m, 4H), 2.20-2.40 (m, 1H), 3.80 (s, 3H), 3.95 (s, 1H, OH), 6.30-6.50 (m, 2H), 7.35-7.45 (m, 1H).

MS: $[\text{M}]^+ = 238$.

(2-Fur-2-yl-2-oxoacetic acid methyl ester was prepared from the commercially available 2-fur-2-yl-2-oxoacetic acid by a standard method).

Intermediate I-12**Preparation of 2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetic acid methyl ester.**

98.6 ml (0.197 mol) of a 2M solution of Cyclohexylmagnesium chloride in diethyl ether, were added to a solution of 27.97 g (0.164 mol) of 2-oxo-2-thien-2-ylacetic acid methyl ester dissolved in 370 ml of THF, at -78°C under a N_2 atmosphere. The mixture was stirred at this temperature for 10 minutes, and then warmed to room temperature. After 1h, the reaction mixture was treated with a saturated solution of ammonium chloride and extracted three times with ethyl acetate. The organic phases were combined, washed with water, and dried over MgSO_4 . After removal of the solvent, the oil obtained was purified by column chromatography (silica gel) using hexane/AcOEt 90:10 as eluent. The yield was 16.2 g (39.5%) of a pure product, whose structure was confirmed by $^1\text{H-NMR}$.

$^1\text{H-NMR}$ (CDCl_3): δ 1.0-1.55 (m, 6H), 1.55-1.90 (m, 4H), 2.0-2.20 (m, 1H), 3.80 (s, 3H), 4.0 (s, 1H, OH), 7.0 (m, 1H), 7.10 (m, 1H), 7.20-7.30 (m, 1H).

MS: $[\text{M}^+]$: 254.

(Intermediate I-12 is described in E. Atkinson et al. J. Med. Chem., (1977), Vol 20, N° 12, 1612-1617)

Intermediate I-13**Preparation of 2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid methyl ester.**

Prepared as in Example I-12. The yield was 3.83 g (37%).

$^1\text{H-NMR}$ (CDCl_3): δ 1.35-1.75 (m, 8H), 2.70-2.90 (m, 1H), 3.80 (s, 3H), 4.02 (s, 1H, OH), 6.95-7.05 (m, 1H), 7.10-7.20 (m, 1H), 7.20-7.25 (m, 1H).

(Intermediate I-13 is described in E. Atkinson et al. J. Med. Chem., (1977), Vol 20, N° 12, 1612-1617)

Method - d -**Intermediate I-14****Preparation of 2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.**

Cyclopentylmagnesium chloride, 0.0472 mol (23.6 ml of a solution 2M in ether) was added to a solution of 9.4 g (0.0377 mol) of 2-fur-2-yl-2-oxoacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester dissolved in 125 ml of THF, at -70°C under a N_2 atmosphere. The mixture was stirred at this temperature for 10 minutes, and then

warmed to room temperature. After 16 h, the reaction mixture was treated with a saturated solution of ammonium chloride and extracted twice with ethyl acetate. The organic phases were combined, washed with water, and dried over MgSO_4 . After removal of the solvent, the oil obtained (7.5 g) was purified by chromatography on silica gel eluting with chloroform/methanol/ammonia 95:5:0.5. Appropriate fractions were combined and evaporated to obtain the two diastereomers: I-14a, I-14b

Intermediate I-14a

(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

1.55 g of an oil corresponding to the first eluted diastereomer was triturated with a mixture of isopropyl ether/ petroleum ether 1:1, to give 0.24 g (4%, based on single isomer) of a solid whose structure was confirmed by $^1\text{H-NMR}$ as a pure diastereomer; mp = 109.6-110.6°C.

$[\alpha]^{22}_{\text{D}} = +19.7^\circ$ (c=1, CHCl_3)

$^1\text{H-NMR}$ (CDCl_3): δ 1.22-1.37 (m, 1H), 1.40-1.80 (m, 11 H), 1.97 (m, 1H), 2.74-2.96 (m, 6 H), 3.19-3.30 (m, 1H), 3.80 (bs, 1H, OH), 4.85-4.89 (m, 1H), 6.34-6.37 (m, 2H), 7.35 (m, 1H)

MS: $[\text{M}+1]^+ = 320$.

Intermediate I-14a was hydrolysed (EtOH/NaOH 2N, 2 h r.t.) to give (+)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid: $[\alpha]^{22}_{\text{D}} = +31.95^\circ$ (c=1, EtOH).

Configuration S was assigned. (See TABLE 1).

Intermediate I-14b

(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

1.10 g of a solid corresponding to the second eluted diastereomer was triturated with a mixture of isopropyl ether/petroleum ether 1:1, to give 0.42 g (7%, based on single isomer) of a solid whose structure was confirmed by $^1\text{H-NMR}$ as a pure diastereomer; mp = 119.9-122.1°C.

$[\alpha]^{22}_{\text{D}} = -14.2^\circ$ (c=1, CHCl_3)

$^1\text{H-NMR}$ (CDCl_3): δ 1.40-1.90 (m, 12H), 2.07 (m, 1H), 2.49-2.56 (m, 1H), 2.67-2.86 (m, 5H), 3.12-3.24 (m, 1H), 3.80 (bs, 1H, OH), 4.87-4.91 (m, 1H), 6.35-6.39 (m, 2H), 7.38 (m, 1H)

MS: $[\text{M}+1]^+ = 320$

Intermediate I-14b was hydrolysed (EtOH/NaOH 2N, 2 h r.t.) to give (-)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid: $[\alpha]^{22}_{\text{D}} = -32.10^\circ$ (c=1, EtOH).

Configuration R was assigned. (See TABLE 1).

(2-Fur-2-yl-2-oxoacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester may be prepared as described in WO 01/04118)

5

Intermediate I-15

Preparation of 2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

Prepared using the same method as for Intermediate I-14, but from from 20.8 g (0.0784 mol) of of 2-oxo-2-thien-2-ylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester and cyclopentylmagnesium chloride, 0.08 mol (40 ml of a solution 2M in ether). The oil obtained (15.64 g) was purified by chromatography on silica gel eluting with chloroform/methanol/ammonia 97:3:0.5 to obtain 8.38 g (32%) of a pure product, mixture of diastereomers: I-15a and I-15b. Structure confirmed by ¹H-NMR.

15 Intermediate I-15a

(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

The oil mixture of diastereomers was treated with isopropyl ether to obtain a solid, that was treated again with isopropyl ether to yield 2.2 g of a pure diastereomer (Intermediate I-15a , 16.7 % based on a single isomer).

$[\alpha]^{22}_D = -5.75^\circ$ (c=1, CHCl₃) ; mp: 152-157°C.

¹H-NMR (CDCl₃): δ 1.40-1.80 (m, 11H), 1.80-2.0 (m, 1H), 2.10 (m, 1H), 2.52-2.65 (m, 1H), 2.70-2.95 (m, 5H), 3.10-3.22 (m, 1H), 4.07 (s, 1H, OH), 4.85-4.95 (m, 1H), 6.95-7.05 (m, 1H), 7.10-7.20 (m, 1H), 7.20-7.27 (m, 1H).

25 MS: [M+1]⁺ = 336

Intermediate I-15a was hydrolysed (EtOH/NaOH 2N, 2 h r.t. and 2h at 60°C) to give (-)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid : $[\alpha]^{22}_D = -6.44^\circ$ (c=1, EtOH).

Configuration S was assigned. (See TABLE 1).

Intermediate I-15b

30 **(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.**

The mother liquors of the first crystallization of intermediate I-15a were evaporated, and the oil obtained treated with isopropyl ether to give a solid, that was treated again with isopropyl ether to yield 1.47 g of the second diastereomer intermediate I-15b (11.2 % based on single isomer).

35 $[\alpha]^{22}_D = +22.49^\circ$ (c=1, CHCl₃) ; mp: 99-102°C

¹H-NMR (CDCl₃): 8.05-1.85 (m, 12H), 2.0 (m, 1H), 2.65-2.95 (m, 6H), 3.22-3.34 (m, 1H), 4.05 (s, 1H, OH), 4.80-4.92 (m, 1H), 6.90-7.0 (m, 1H), 7.00-7.16 (m, 1H), 7.20-7.27 (m, 1H).

MS: [M+1]⁺ = 336

- 5 Intermediate I-15b was hydrolysed (EtOH/NaOH 2N, 2 h r.t. and 2 h at 60°C) to give (+)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid : $[\alpha]_{D}^{22} = + 6.63^{\circ}$ (c=1, EtOH). Configuration R was assigned. (See TABLE 1).

(Intermediates I-15a and I-15b have also been prepared following method c)

10

(2-Oxo-2-thien-2-ylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester may be prepared as described in WO 01/04118)

Intermediate I-16

- 15 **Preparation of 2-Cyclohexyl-2-hydroxy-2-phenylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.**

Cyclohexylmagnesium chloride, 0.127 mol (63.6 ml of a solution 2M in ether), was added to a solution of 28.7 g (0.111 mol) of 2-oxo-2-phenylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester dissolved in 350 ml of THF, at -70°C under a N₂ atmosphere. The mixture was stirred at this temperature for 10 minutes, and then warmed to room temperature. After 1h, the reaction mixture was treated with a saturated solution of ammonium chloride and extracted twice with ethyl acetate. The organic phases were combined, washed with water, and dried over MgSO₄. After removal of the solvent, the oil obtained (27.0 g) was purified by chromatography on silica gel eluting with chloroform/methanol 10:1. The yield was 18.7 g (49.2%) of a pure product, mixture of diastereomers: I-16a and I-16b.

25

Intermediate I-16a

(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester

- 30 16 g of the oil mixture of diastereomers were dissolved in isopropanol and treated with 5.4 g of fumaric acid. After cooling to 0-5°C, 8 g of the fumarate salt of the first diastereomer (Intermediate I-16a) were obtained.

Fumarate salt: MS: [M (free base) + 1]⁺ = 344.

- The 8 g of this salt were recrystallised from isopropanol to obtain 5 g of a more pure product. This salt was treated with CHCl₃ and K₂CO₃ solution to obtain the free base Intermediate I-16a.

35

I-16a (free base):

$[\alpha]^{22}_D = -14.9^\circ$ ($c=1$, CHCl_3).

$^1\text{H-NMR}$ (CDCl_3): δ 1.0-1.95 (m, 14H), 2.04-2.12 (m, 1H), 2.16-2.32 (m, 1H), 2.38-2.50 (m, 1H), 2.64-2.96 (m, 4H), 3.04-3.16 (m, 1H), 3.70-3.85 (s, 1H, OH), 4.85-4.90 (m, 1H), 7.25-7.40 (m, 3H), 7.60-7.70 (m, 2H).

- 5 Intermediate I-16a (free base) was hydrolysed (EtOH/NaOH 2N, 7 h at 60°C) to give (-)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid as a pure enantiomer $[\alpha]^{22}_D = -23.6^\circ$ ($c=1.4$, EtOH). This value was assigned to the R configuration provided that in the literature (A. Tambuté, A. Collet; Bulletin de la Société Chimique de France, 1984, N° 1-2, pages II-77 to II-82) the (2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid has
10 been described with $[\alpha]^{24}_D = +25.2^\circ$ ($c=1.4$, EtOH). (See TABLE 1).

Intermediate I-16b

(2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester

- The mother liquors of the fumarate salt of the Intermediate I-16a were evaporated and
15 the residue dissolved in water. The obtained solution was washed with ether, basified with K_2CO_3 and extracted with CHCl_3 . The organic layer was dried over MgSO_4 and the solvent was evaporated. The oil obtained (7.5 g) was dissolved in 50 ml of isopropanol and treated with EtOH/HCl(g) . After the addition of 75 ml of ethyl ether, 3.1 g of the chlorhydrate salt of the second diastereomer, Intermediate I-16b, were obtained.

- 20 Chlorhydrate salt: MS: $[\text{M}(\text{free base}) + 1]^+ = 344$.

The chlorhydrate salt was treated with CHCl_3 and K_2CO_3 solution to obtain the free base Intermediate I-16b.

I-16b (free base):

$[\alpha]^{22}_D = +25.3^\circ$ ($c=1$, CHCl_3).

- 25 $^1\text{H-NMR}$ (CDCl_3): δ 1.0-1.78 (m, 13H), 1.78-1.90 (m, 1H), 1.92-2.0 (m, 1H), 2.20-2.34 (m, 1H), 2.66-2.96 (m, 5H), 3.20-3.32 (m, 1H), 3.70-3.85 (s, 1H, OH), 4.75-4.85 (m, 1H), 7.25-7.40 (m, 3H), 7.60-7.70 (m, 2H).

- Intermediate I-16b (free base) was hydrolysed (EtOH/NaOH 2N, 60°C , 8 h) to give (+)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid as a pure enantiomer $[\alpha]^{22}_D = +23.1^\circ$
30 ($c=1.4$, EtOH). This value was assigned to the S configuration according with the results obtained with Intermediate I-16a. (See TABLE 1).

(2-Oxo-2-phenylacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester may be prepared as described in WO 92/04346.)

Intermediate I-17**2-hydroxy-2,3-diphenylpropionic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester**

Prepared using the same method as for Intermediate I-16 as a mixture of
5 diastereomers I-17a and I-17b, which were separated by crystallisation using
ether/isopropyl ether.

Intermediate I-17a

(2*)-2-hydroxy-2,3-diphenylpropionic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester,
first diastereomer obtained.

10 The yield was 0.87 g (42.6% based on single isomer), mp: 132°C.

¹H-NMR (CDCl₃): δ 1.30-1.60 (m, 2H), 1.60-1.90 (m, 2H), 2.05 (m, 1H), 2.20-2.35 (m,
1H), 2.50-2.90 (m, 4H), 3.0-3.15 (m, 1H), 3.25 and 3.60 (dd, 2H), 3.70 (bs, 1H, OH),
4.70-4.80 (m, 1H), 7.15-7.45 (m, 8H), 7.65-7.75 (m, 2H).

MS: [M+1]⁺ = 352

15 ((*): Configuration not assigned)

Intermediate I-17b

(2*)-2-hydroxy-2,3-diphenylpropionic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester,
second diastereomer obtained

The yield was 0.23 g (11.2% based on single isomer), mp: 107°C.

20 ¹H-NMR (CDCl₃): δ 1.20-1.35 (m, 1H), 1.35-1.55 (m, 2H), 1.55-1.70 (m, 1H), 1.80-1.95
(m, 1H), 2.55-2.90 (m, 5H), 3.10-3.20 (m, 1H), 3.25 and 3.60 (dd, 2H), 3.80 (bs, 1H,
OH), 4.65-4.80 (m, 1H), 7.20-7.50 (m, 8H), 7.65-7.75 (m, 2H).

MS: [M+1]⁺ = 352

((*): Configuration not assigned)

25

Intermediate I-18

**2-Hydroxy-3-phenyl-2-thien-2-ylpropionic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl
ester.**

Prepared using the same method as for Intermediate I-15 as a mixture of
30 diastereomers. The yield was 0.81 g (54 %). The product was purified by preparation
of the chlorhydrate salt, 0.57 g of this salt were obtained (63 % from the free base).

Chlorhydrate salt:

¹H-NMR (DMSO-d₆): δ 1.40-1.60 (m, 1H), 1.60-1.95 (m, 3H), 2.05 and 2.10 (m, 1H),
2.75-3.65 (m, 8H), 4.90-5.05 (m, 1H), 6.50 and 6.55 (s, 1H, OH), 6.95-7.05 (m, 1H),
35 7.10-7.30 (m, 6H), 7.40-7.50 (m, 1H), 10.9 (bs, 1H, NH⁺).

MS: [M+1]⁺ = 358

Intermediate I-19**2-Hydroxy-2-thien-2-ylpent-3-ynoic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.**

Prepared using the same method as for Intermediate I-15 as a mixture of diastereomers. The yield was 1.88 g (25.6%)

¹H-NMR (CDCl₃): δ 1.20-1.90 (m, 4H), 1.92 and 1.96 (s, 3H, CH₃), 2.0 and 2.16 (m, 1H), 2.45-2.90 (m, 5H), 3.05-3.20 and 3.15-3.27 (m, 1H), 4.85-4.92 (m, 1H), 6.94-7.0 (m, 1H), 7.24-7.30 (m, 2H), the signal for OH group is observed between 4.5 and 5.5 as a broad band.

MS: [M+1]⁺ = 306

Intermediate I-20**2-Hydroxy-2-thien-2-ylbut-3-enoic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.**

Prepared using the same method as for Intermediate I-15 as a mixture of diastereomers. The yield was 1.74 g (18.8%)

¹H-NMR (CDCl₃): δ 1.30-1.90 (m, 4H), 2.05-2.15 (m, 1H), 2.60-3.0 (m, 5H), 3.15-3.35 (m, 1H), 3.40-4.70 (broad band, 1H, OH), 4.85-4.95 (m, 1H), 5.30-5.40 (m, 1H), 5.60-5.75 (m, 1H), 6.30-6.50 (m, 1H), 6.95-7.05 (1H), 7.10-7.15 (m, 1H), 7.25-7.30 (m, 1H).

MS: [M+1]⁺ = 294

Intermediate I-21**2-Hydroxy-4-phenyl-2-thien-2-ylbutyric acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.**

Prepared using the same method as for Intermediate I-15 as a mixture of diastereomers. The yield was 0.29 g (2.4%).

¹H-NMR (CDCl₃): δ 1.25-1.95 (m, 4H), 1.95-2.10 (m, 1H), 2.30-3.0 (m, 9H), 3.10-3.25 (m, 1H), 4.80-4.90 (m, 1H), 6.95-7.05 (m, 1H), 7.05-7.40 (m, 7H).

MS: [M+1]⁺ = 372

Method - e -**Intermediate I-22**

Preparation of 2-Hydroxymethyl-2,3-diphenylpropionic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester

Lithium diisopropylamide (0.0048 mol) 2.40 ml of a 2M solution (in heptane/THF/ethylbenzene) was added to a stirred solution of 1.5 g (0.0045 mol) of

2,3-diphenylpropionic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester in 30 ml of THF at -70°C under a N_2 atmosphere. CH_2O (gas) was bubbled into the reaction mixture via a steady stream of dry N_2 during 10 min at -70°C and then while the mixture was warmed at room temperature. The reaction was quenched by addition of saturated ammonium chloride solution (100 ml) and the resultant mixture was extracted twice with 100 ml of ethyl acetate. The organic layers were combined, dried over MgSO_4 and evaporated to yield 1.9 g of an oil. This 1.9 g were combined with 3.28 g of a previous preparation and the product obtained (5.18 g) was purified by chromatography on silica gel eluting with chloroform/methanol/ammonia from 97.5:2.5:0.25 to 90:10:1. Appropriate fractions were combined and evaporated to obtain the two diastereomers : Intermediates I-22a and I-22b

Intermediate I-22a :

(2*)-2-Hydroxymethyl-2,3-diphenylpropionic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

1.25 g of a solid corresponding to the first eluted diastereomer was triturated with isopropyl ether, to give 0.95 g (42%, based on single isomer) of a white solid whose structure was confirmed by ^1H -NMR as a pure diastereomer ; mp: 119°C .

^1H -NMR (CDCl_3): δ 1.20-1.35 (m, 1H), 1.40-1.70 (m, 3H), 1.90 (m, 1H), 2.5 (bs, OH), 2.60-2.85 (m, 5H), 3.15-3.25 (m, 1H), 3.40-3.50 (dd, 2H), 3.95-4.10 (dd, 2H), 4.85 (m, 1H), 7.05 (m, 2H), 7.15-7.40 (m, 8H)

MS: $[\text{M}+1]^+ = 366$

((*): Configuration not assigned)

Intermediate I-22b

(2*)-2-Hydroxymethyl-2,3-diphenylpropionic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester.

1.84 g of an oil corresponding to the second eluted diastereomer was purified by column chromatography (in the above described conditions) to yield 1.26 g of a solid that after trituration with isopropyl ether yield 0.95 g of a white solid whose structure was confirmed by ^1H -NMR as a pure diastereomer (42%, based on single isomer); mp: 154°C .

^1H -NMR (CDCl_3): δ 1.20-1.35 (m, 1H), 1.50-1.75 (m, 3H), 2.0 (m, 1H), 2.35 (bs, OH), 2.50-2.80 (m, 5H), 3.10-3.20 (m, 1H), 3.35-3.50 (dd, 2H), 3.95-4.10 (dd, 2H), 4.85 (m, 1H), 7.0 (m, 2H), 7.15-7.40 (m, 8H).

MS: $[\text{M}+1]^+ = 366$

((*): Configuration not assigned)

Also included within the scope of the present invention are pharmaceutical compositions which comprise, as the active ingredient, at least one quinuclidine derivative of general formula (I) in association with a pharmaceutically acceptable carrier or diluent. Preferably the composition is made up in a form suitable for oral administration.

The pharmaceutically acceptable carrier or diluents which are mixed with the active compound or compounds, to form the composition of this invention are well-known *per se* and the actual excipients used depend *inter alia* on the intended method of administration of the composition.

Compositions of this invention are preferably adapted for oral administration. In this case, the composition for oral administration may take the form of tablets, film-coated tablets, liquid inhalant, powder inhalant and inhalation aerosol; all containing one or more compounds of the invention; such preparations may be made by methods well-known in the art.

The diluents which may be used in the preparations of the compositions include those liquid and solid diluents which are compatible with the active ingredient, together with colouring or flavouring agents, if desired. Tablets or film-coated tablets may conveniently contain between 1 and 500 mg, preferably from 5 to 300 mg of active ingredient. The inhalant compositions may contain between 1 µg and 1,000 µg, preferably from 10 µg to 800 µg of active ingredient. In human therapy, the dose of the compound of general formula (I) depend on the desired effect and duration of treatment; adult doses are generally between 3 mg and 300 mg per day as tablets and 10 µg and 800 µg per day as inhalant composition.

Pharmacological Action

The results on human muscarinic receptors binding and in the test on bronchospasm in guinea pig, were obtained as described below.

Human muscarinic receptor studies.

The binding of [3H]-NMS to human muscarinic receptors was performed according to Waelbroek et al (1990), Mol. Pharmacol., 38: 267-273. Assays were carried out at 25°C. Membrane preparations from stably transfected chinese hamster ovary-K1 cells (CHO) expressing the genes for the human muscarinic receptors M3 were used.

For determination of IC_{50} , membrane preparations were suspended in DPBS to a final concentration of 89 $\mu\text{g/ml}$ for the M3 subtype. The membrane suspension was incubated with the tritiated compound for 60 min. After incubation the membrane fraction was separated by filtration and the bound radioactivity determined. Non specific binding was determined by addition of 10^{-4} M atropine. At least six concentrations were assayed in duplicate to generate individual displacement curves.

Our results show that the compounds of the present invention have high affinities for muscarinic M3 receptors, preferably human muscarinic receptors. Affinity levels have been measured by in vitro assays as described above at 100 nM and 10 nM. Preferred compounds of the invention produce an inhibition of $[^3\text{H}]\text{-NMS}$ binding of at least 35 % at 10 nM and of at least 65 % at 100 nM (Table 2).

TABLE 2

N° COMPOUND	% INHIBITION concentration: 1,00E-07 M	% INHIBITION concentration: 1,00E-08 M
Atropine	88,3	69,75
Ipratropium bromide	93,75	67,25
13	76	36
14	76.5	39.0
16	74.2	36.3
22	81.5	72
23	75.6	63.3
24	78	56.6
25	76.1	62.6
26	75.6	63.8
28	78.3	60.6
29	79.0	53.8
31	74.3	54.3
32	73.9	44.5
33	72.8	46.7
34	85.3	68.3
36	84.2	42.0
37	88.1	72.6
38	86.3	57.7
40	86.9	72.7
43	83.4	58.7
44	84.6	44.6
46	87.1	57.6
53	81.5	58
54	72.5	44.1
56	77.3	53.8
57	77.4	47.1
61	75.1	39.9
64	78.6	64.5
65	79.8	66.0
67	75.1	52.5
69	70.8	43.9
70	71.2	50.0
80	72.4	55.8
81	70.1	45.4
82	70.6	55.3
83	72.7	60.3
84	68.3	41.0
86	68.2	37.2
88	65.5	35.7
89	68.5	51.3
92	69.4	49.2

Test on bronchospasm in guinea pig

The studies were performed according to H. Konzett and F. Rössler (1940), Arch. Exp. Path. Pharmacol. 195: 71-74. Aqueous solutions of the agents to be tested were nebulized and inhaled by anaesthetized ventilated male guinea pigs (Dunkin-Hartley). Bronchial response to intravenous acetylcholine challenge was determined before and after drug administration and the changes in pulmonary resistance at several time-points were expressed as percent of inhibition of bronchoespasm.

The compounds of the present invention inhibited the bronchospasm response to acetylcholine with high potency and a long duration of action.

From the above described results one of ordinary skill in the art can readily understand that the compounds of the present invention have excellent antimuscarinic activity (M3) and thus are useful for the treatment of diseases in which the muscarinic M3 receptor is implicated, including respiratory disorders such as chronic obstructive pulmonary disease (COPD), bronchitis, bronchial hyperreactivity, asthma, cough and rhinitis; urological disorders such as urinary incontinence, pollakiuria, neurogenic or unstable bladder, cystospasm and chronic cystitis; gastrointestinal disorders such as irritable bowel syndrome; spastic colitis, diverticulitis and peptic ulceration; and cardiovascular disorders such as vagally induced sinus bradycardia. For example, the compounds of the present invention are useful for the treatment of respiratory diseases such as chronic obstructive pulmonary disease, chronic bronchitis, asthma, and rhinitis; urinary diseases such as urinary incontinence and pollakinuria in neuripenia pollakinuria, neurogenic bladder, nocturnal enuresis, unstable bladder, cytospsasm and chronic cystitis; and gastrointestinal diseases such as irritable bowel syndrome, spastic colitis and diverticulitis.

The present invention further provides a compound of formula (I) or a pharmaceutically acceptable composition comprising a compound of formula (I) for use in a method of treatment of the human or animal body by therapy, in particular for the treatment of respiratory, urological or gastrointestinal disease or disorder.

The present invention further provides the use of a compound of formula (I) or a pharmaceutically acceptable composition comprising a compound of formula (I) for the manufacture of a medicament for the treatment of a respiratory, urological or gastrointestinal disease or disorder.

Further, the compounds of formula (I) and pharmaceutical compositions comprising a compound of formula (I) can be used in a method of treating a respiratory, urological or gastrointestinal disease or disorder, which method comprises

administering to a human or animal patient in need of such treatment an effective, non-toxic, amount of a compound of formula (I) or a pharmaceutical composition comprising a compound of formula (I).

Further, the compounds of formula (I) and pharmaceutical compositions comprising a compound of formula (I) can be used in combination with other drugs effective in the treatment of these diseases. For example with β_2 agonists, steroids, antiallergic drugs, phosphodiesterase IV inhibitors and/or leukotriene D4 (LTD4) inhibitors, for simultaneous, separate or sequential use in the treatment of a respiratory disease.

The present invention therefore further provides a combination product comprising

- (i) a compound according to the invention; and
- (ii) another compound effective in the treatment of a respiratory, urological or gastrointestinal disease or disorder

for simultaneous, separate or sequential use.

The compound (ii) which is effective in the treatment of a respiratory, urological or gastrointestinal disease or disorder may be a β_2 agonist, steroid, antiallergic drug, phosphodiesterase IV inhibitor and/or leukotriene D4 (LTD4) antagonist when the product is for simultaneous, separate or sequential use in the treatment of a respiratory disease. Alternatively, the compound (ii) may be a β_2 agonist, steroid, antiallergic drug and/or phosphodiesterase IV inhibitor when the product is for simultaneous, separate or sequential use in the treatment of a respiratory disease.

The present invention will be further illustrated by the following examples. The examples are given by way of illustration only and are not to be construed as limiting.

Example 1

(3R)-3-(2,3-Diphenylpropionyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate.

The title compound was synthesised as a mixture of diastereomers according to methods c, and b from the Intermediate I-2. The yield of the final step was 20 mg, 71%.

MS $[M-CF_3COO]^+$: 470.

1H -NMR (DMSO- d_6): δ 1.50-1.72 (m, 2H), 1.75-1.95 (m, 2H), 1.97-2.15 (m, 3H), 2.95-3.15 (m, 4H), 3.20-3.50 (m, 5H), 3.75-3.85 (m, 1H), 3.95-4.15 (m, 3H), 4.95-5.05 (m, 1H), 6.90-7.0 (m, 4H), 7.15-7.45 (m, 11H).

Example 2**(3R)-3-(2,3-Diphenylpropionyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

5 The title compound was synthesised as a mixture of diastereomers according to methods c, and b from the Intermediate I-2. The yield of the final step was 15 mg, 55%.

MS [M-CF₃COO]⁺: 460.

¹H-NMR (DMSO-d₆): δ 1.50-1.70 (m, 2H), 1.70-2.0 (m, 4H), 2.0-2.15 (m, 1H), 2.75-2.85 (m, 2H), 2.85-3.20 (m, 4H), 3.20-3.45 (m, 5H), 3.70-3.82 (m, 1H), 4.02-4.12 (m, 1H),
10 4.95-5.02 (m, 1H), 6.90-7.05 (m, 2H), 7.10-7.45 (m, 11H)

Example 3**(3R)-3-[(2*)-2-Hydroxymethyl-2,3-diphenylpropionyloxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 1)**

15 The title compound was synthesised according to methods e, and b from the Intermediate I-22a. The yield of the final step was 15 mg, 52%.

MS [M-CF₃COO]⁺: 490.

¹H-NMR (DMSO-d₆): δ 1.55-1.75 (m, 2H), 1.75-2.05 (m, 4H), 2.21 (m, 1H), 2.75-2.85 (m, 2H), 2.85-2.95 (m, 1H), 3.05-3.45 (m, 8H), 3.75-3.87 (m, 2H), 3.92-4.0 (m, 1H),
20 5.08 (m, 1H), 5.20-5.23 (t, 1H, OH), 6.82-6.90 (m, 2H), 6.90-6.95 (m, 1H), 6.95-7.02 (m, 1H), 7.05-7.20 (m, 5H), 7.20-7.35 (m, 3H), 7.37-7.42 (m, 1H).

Example 4**(3R)-3-[(2*)-2-Hydroxymethyl-2,3-diphenylpropionyloxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 1)**

25 The title compound was synthesised according to methods e, and b from the Intermediate I-22a. The yield of the final step was 18 mg, 64%.

MS [M-CF₃COO]⁺: 470.

¹H-NMR (DMSO-d₆): δ 1.62-1.75 (m, 2H), 1.80-2.05 (m, 2H), 2.26 (m, 1H), 2.90-3.12 (m, 3H), 3.20-3.55 (m, 8H), 3.80-4.02 (m, 3H), 5.10-5.17 (m, 1H), 5.20-5.25 (t, 1H, OH); 6.82-6.90 (m, 2H), 7.10-7.20 (m, 5H), 7.22-7.40 (m, 8H).

Example 5**(3R)-3-[(2*)-2-Hydroxymethyl-2,3-diphenylpropionyloxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 2)**

The title compound was synthesised according to methods e, and b from the Intermediate I-22b. The yield of the final step was 10.1 mg, 37%.

MS [M-CF₃COO]⁺: 490.

¹H-NMR (DMSO-d₆): δ 1.45-1.60 (m, 1H), 1.60-1.75 (m, 1H), 1.80-2.05 (m, 4H), 2.18 (m, 1H), 2.75-2.90 (m, 2H), 2.95-3.10 (m, 1H), 3.10-3.55 (m, 8H), 3.75-3.92 (m, 2H), 4.0-4.12 (m, 1H), 5.05-5.15 (m, 1H), 5.25-5.35 (t, 1H, OH), 6.70-6.85 (m, 2H), 6.90-7.20 (m, 7H), 7.20-7.35 (m, 3H), 7.35-7.42 (m, 1H).

10 Example 6

(3R)-3-[(2*)-2-Hydroxymethyl-2,3-diphenylpropionyloxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 2)

The title compound was synthesised according to methods e, and b from the Intermediate I-22b. The yield of the final step was 22 mg, 76%.

15 MS [M-CF₃COO]⁺: 470.

¹H-NMR (DMSO-d₆): δ 1.50-1.60 (m, 1H), 1.60-1.80 (m, 1H), 1.85-2.05 (m, 2H), 2.21 (m, 1H), 2.90-3.10 (m, 2H), 3.12-3.55 (m, 9H), 3.78-3.83 (m, 1H), 3.88-3.95 (m, 1H), 4.07-4.12 (m, 1H), 5.15-5.20 (m, 1H), 5.35-5.40 (t, 1H, OH), 6.75-6.80 (m, 2H), 7.0-7.15 (m, 3H), 7.20-7.40 (m, 8H).

20

Example 7

(3R)-3-[(2*)-2-Hydroxy-2,3-diphenylpropionyloxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide (diastereomer 1)

25 The title compound was synthesised according to methods d, and a from the Intermediate I-17a. The yield of the final step was 380 mg, 96%.

MS [M-Br]⁺: 486, mp: 103°C.

¹H-NMR (DMSO-d₆): δ 1.65-1.75 (m, 2H), 1.75-2.02 (m, 2H), 2.02-2.15 (m, 2H), 2.24 (m, 1H), 3.05-3.25 (m, 2H), 3.25-3.55 (m, 7H), 3.78-3.90 (m, 1H), 3.98-4.08 (m, 2H), 30 5.02-5.10 (m, 1H), 6.20 (s, 1H, OH), 6.92-7.0 (m, 3H), 7.10-7.22 (m, 5H), 7.25-7.40 (m, 5H), 7.52-7.58 (m, 2H).

Example 8

(3R)-3-[(2*)-2-Hydroxy-2,3-diphenylpropionyloxy]-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide (diastereomer 1)

35

The title compound was synthesised according to methods d, and a from the Intermediate I-17a. The yield of the final step was 320 mg, 83%.

MS [M-Br]⁺: 472, m/z 223°C.

¹H-NMR (DMSO-d₆): δ 1.70-1.80 (m, 2H), 1.80-2.0 (m, 2H), 2.2 (m, 1H), 3.15-3.55 (m, 7H), 3.55-3.70 (m, 2H), 3.85-4.0 (m, 1H), 4.30-4.45 (m, 2H), 5.0-5.10 (m, 1H), 6.10 (s, 1H, OH), 6.90-7.05 (m, 3H), 7.05-7.20 (m, 5H), 7.20-7.40 (m, 5H), 7.45-7.55 (m, 2H).

5

Example 9

(3R)-3-[(2*)-2-Hydroxy-2,3-diphenylpropionyloxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 2)

The title compound was synthesised according to methods d, and b from the Intermediate I-17b. The yield of the final step was 7.2 mg, 25%.

10

MS [M-CF₃COO]⁺: 486.

¹H-NMR (DMSO-d₆): δ 1.40-1.55 (m, 1H), 1.55-1.70 (m, 1H), 1.75-2.0 (m, 2H), 2.11 (m, 3H), 3.10-3.60 (m, 9H), 3.77-3.87 (m, 1H), 4.0-4.1 (m, 2H), 5.0-5.1 (m, 1H), 6.14 (s, 1H, OH), 6.90-7.0 (m, 3H), 7.15-7.25 (m, 5H), 7.25-7.42 (m, 5H), 7.60-7.67 (m, 2H).

15

Example 10

(3R)-3-[(2*)-2-Hydroxy-2,3-diphenylpropionyloxy]-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate (diastereomer 2)

The title compound was synthesised according to methods d, and b from the Intermediate I-17b. The yield of the final step was 5.4 mg, 19%.

20

MS [M-CF₃COO]⁺: 472.

¹H-NMR (DMSO-d₆): δ 1.40-1.55 (m, 1H), 1.55-1.70 (m, 1H), 1.80-2.0 (m, 2H), 2.12 (m, 1H), 3.20-3.60 (m, 7H), 3.60-3.70 (m, 2H), 3.90-4.0 (m, 1H), 4.42 (m, 2H), 5.0-5.1 (m, 1H), 6.15 (s, 1H, OH), 6.95-7.05 (m, 3H), 7.10-7.22 (m, 5H), 7.25-7.40 (m, 5H), 7.57-7.65 (m, 2H).

25

Example 11

(3R)-3-(2-Hydroxy-3-phenyl-2-thien-2-ylpropionyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

30

The title compound was synthesised as a mixture of diastereomers according to methods d, and b from Intermediate I-18. The yield of the final step was 15 mg, 52%.

MS [M-CF₃COO]⁺: 492.

¹H-NMR (DMSO-d₆): δ 1.45-1.70 (m, 1H), 1.75-2.0 (m, 3H), 2.0-2.30 (m, 3H), 3.0-3.17 (m, 1H), 3.17-3.57 (m, 8H), 3.80-3.90 (m, 1H), 3.97-4.10 (m, 2H), 5.02-5.05 (m, 1H),

35

6.52-6.60 (d, 1H, OH), 6.90-7.04 (m, 4H), 7.14-7.28 (m, 6H), 7.28-7.38 (m, 2H), 7.42-7.50 (m, 1H).

5 **Example 12**

(3R)-3-(2-Hydroxy-3-phenyl-2-thien-2-ylpropionyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as a mixture of diastereomers according to methods d, and b from Intermediate I-18. The yield of the final step was 21 mg, 74%.

10 MS [M-CF₃COO]⁺: 482

¹H-NMR (DMSO-d₆): δ 1.45-1.70 (m, 1H), 1.75-2.05 (m, 5H), 2.05-2.3 (m, 1H), 2.77-2.87 (m, 2H), 2.90-3.10 (m, 1H), 3.10-3.52 (m, 8H), 3.75-3.82 (m, 1H), 5.0-5.07 (m, 1H), 6.52-6.57 (d, 1H, OH), 6.92-7.05 (m, 3H), 7.10-7.27 (m, 6H), 7.37-7.47 (m, 2H).

15

Example 13

(3R)-3-(2-Hydroxy-2-thien-2-ylpent-4-enoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide

20 The title compound was obtained as a mixture of diastereomers according to methods c, and a from Intermediate I-3. The yield of the final step was 300 mg, 71%.

MS [M-Br]⁺: 442, mp: 157°, (described in experimental section, method a)

¹H-NMR (DMSO-d₆): δ 1.70-2.05 (m, 4H), 2.05-2.35 (m, 3H), 2.70-2.83 (m, 1H), 2.90-3.02 (m, 1H), 3.25-3.60 (m, 7H), 3.82-3.97 (m, 1H), 3.97-4.10 (m, 2H), 5.05-5.25 (m, 3H), 5.70-5.90 (m, 1H), 6.50 (d, 1H, OH), 6.90-7.05 (m, 4H), 7.10-7.20 (m, 1H), 7.27-7.35 (m, 2H), 7.45 (m, 1H).

25

Example 14

30 **(3R)-3-(2-Hydroxy-2-thien-2-ylpent-4-enoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was obtained as a mixture of diastereomers according to methods c, and b from Intermediate I-3. The yield of the final step was 10 mg, 39.3%.

MS [M-CF₃COO]⁺: 432.

¹H-NMR (DMSO-d₆): δ 1.65-2.18 (m, 6H), 2.18-2.30 (m, 1H), 2.70-3.05 (m, 4H), 3.10-3.55 (m, 4H), 3.55-3.68 (m, 1H), 3.78-3.92 (m, 2H), 4.0-4.1 (m, 1H), 5.0-5.20 (m, 3H), 5.70-5.85 (m, 1H), 6.48-6.52 (d, 1H, OH), 6.90-7.02 (m, 3H), 7.10-7.20 (m, 1H), 7.35-7.42 (m, 1H), 7.42-7.50 (m, 1H).

35

Example 15**(3R)-3-(2-Hydroxy-2-thien-2-ylpent-4-enoyloxy)-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide**

5 The title compound was obtained as a mixture of diastereomers according to methods c, and a from Intermediate I-3. The yield of the final step was 270 mg , 66 %.

MS [M-Br]⁺: 428, mp: 82°C.

¹H-NMR (DMSO-d₆): δ 1.72-2.10 (m, 4H), 2.20-2.35 (m, 1H), 2.70-2.85 (m, 1H), 2.90-3.05 (m, 1H), 3.25-3.85 (m, 7H), 3.92-4.12 (m, 1H), 4.35-4.45 (m, 2H), 4.95-5.20 (m, 3H), 5.70-5.90 (m, 1H), 6.50 (s, 1H, OH), 6.90-7.05 (m, 4H), 7.10-7.18 (m, 1H), 7.25-7.45 (m, 3H).

Example 16**(3R)-3-(2-Hydroxy-2-thien-2-ylheptanoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was obtained as a mixture of diastereomers according to methods c, and b from Intermediate I-1. The yield of the final step was 16.2 mg , 57 %.

MS [M-CF₃COO]⁺: 472.

20 ¹H-NMR (DMSO-d₆): δ 0.80-0.90 (m, 3H), 1.15-1.40 (m, 6H), 1.65-2.30 (m, 9H), 3.20-3.60 (m, 5H), 3.85-3.95 (m, 1H), 3.95-4.10 (m, 2H), 5.05-5.17 (m, 1H), 6.30-6.35 (d, 1H, OH), 6.90-7.05 (m, 4H), 7.10-7.17 (m, 1H), 7.25-7.35 (m, 2H), 7.42-7.48 (m, 1H).

Example 17**(3R)-3-(2-Hydroxy-2-thien-2-ylheptanoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate.**

The title compound was obtained as a mixture of diastereomers according to methods c, and b from Intermediate I-1. The yield of the final step was 6.8 mg , 12 %.

30 MS [M-CF₃COO]⁺: 462.

¹H-NMR (DMSO-d₆): δ 0.80-0.90 (m, 3H), 1.15-1.40 (m, 6H), 1.65-2.30 (m, 9H), 2.80-2.85 (m, 2H), 3.10-3.55 (m, 7H), 3.75-3.90 (m, 1H), 5.10 (m, 1H), 6.30-6.32 (d, 1H, OH), 6.90-6.95 (m, 1H), 6.95-7.02 (m, 2H), 7.09-7.13 (m, 1H), 7.37-7.39 (m, 1H), 7.40-7.45 (m, 1H).

35

Example 18

(3R)-3-(2-Hydroxy-2-thien-2-ylpent-3-ynoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was obtained as a mixture of diastereomers according to methods d, and b from Intermediate I-19. The yield of the final step was 6.4 mg , 12 %.

MS [M-CF₃COO]⁺: 440.

¹H-NMR (DMSO-d₆): δ 1.60-2.05 (m, 7H), 2.05-2.20 (m, 2H), 2.20-2.35 (m, 1H), 3.10-3.60 (m, 7H), 3.82-3.97 (m, 1H), 3.97-4.10 (m, 2H), 5.13 (m, 1H), 6.90-7.06 (m, 4H), 7.20-7.38 (m, 4H), 7.50-7.56 (m, 1H).

10

Example 19

(3R)-3-(2-Hydroxy-2-thien-2-ylpent-3-ynoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was obtained as a mixture of diastereomers according to methods d, and b from Intermediate I-19. The yield of the final step was 2.4 mg , 4.3 %.

MS [M-CF₃COO]⁺: 430

¹H-NMR (DMSO-d₆): δ 1.60-2.10 (m, 9H), 2.20-2.35 (m, 1H), 2.75-2.90 (m, 2H), 3.10-3.70 (m, 7H), 3.75-3.95 (m, 1H), 5.12 (m, 1H), 6.91-7.04 (m, 3H), 7.19-7.42 (m, 3H), 7.48-7.55 (m, 1H).

20

Example 20

(3R)-3-(2-Hydroxy-2-thien-2-ylbut-3-enoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was obtained as a mixture of diastereomers according to methods d, and b from Intermediate I-20. The yield of the final step was 9.6 mg , 16 %.

MS [M-CF₃COO]⁺: 428

¹H-NMR (DMSO-d₆): δ 1.60-2.05 (m, 4H), 2.05-2.20 (m, 2H), 2.20-2.38 (m, 1H), 3.15-3.60 (m, 7H), 3.82-3.95 (m, 1H), 3.98-4.10 (m, 2H), 5.10-5.20 (m, 1H), 5.25-5.35 (m, 1H), 5.45-5.55 (m, 1H), 6.45-6.55 (m, 1H), 6.75-6.82 (d, 1H, OH), 6.92-6.96 (m, 3H), 6.98-7.03 (m, 1H), 7.13-7.15 (m, 1H), 7.28-7.34 (m, 1H), 7.48-7.52 (m, 1H).

30

Example 21

(3R)-3-(2-Hydroxy-2-thien-2-ylbut-3-enoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

35

The title compound was obtained as a mixture of diastereomers according to methods d, and b from Intermediate I-20. The yield of the final step was 5.8 mg , 10 %.

MS [M-CF₃COO]⁺: 470

¹H-NMR (DMSO-d₆): δ 1.60-2.20 (m, 6H), 2.20-2.35 (m, 1H), 2.84 (m, 2H), 3.10-3.55 (m, 7H), 3.80-3.90 (m, 1H), 5.10-5.20 (m, 1H), 5.25-5.35 (m, 1H), 5.45-5.55 (m, 1H), 6.45-6.55 (m, 1H), 6.75-6.78 (d, 1H, OH), 6.92-6.95 (m, 1H), 6.95-7.05 (m, 2H),
5 7.10-7.15 (m, 1H), 7.35-7.42 (m, 1H), 7.45-7.52 (m, 1H).

Example 22

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide

10 The title compound was synthesised according to methods d, and a from the Intermediate I-15a. The yield of the final step was 230 mg, 85%.

MS [M-Br]⁺: 470, mp: 171°C.

¹H-NMR (DMSO-d₆): δ 1.27-1.67 (m, 8H), 1.8-2.05 (m, 4H), 2.05-2.20 (m, 2H), 2.25-2.35 (m, 1H), 2.70-2.92 (m, 1H), 3.20-3.25 (m, 1H), 3.25-3.60 (m, 6H), 3.80-3.95 (m,
15 1H), 3.95-4.08 (m, 2H), 5.10-5.20 (m, 1H), 6.18 (s, 1H, OH), 6.87-7.05 (m, 4H), 7.08-7.20 (m, 1H), 7.25-7.37 (m, 2H), 7.40-7.47 (m, 1H).

Example 23

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

20 The title compound was synthesised according to methods d, and b from the Intermediate I-15a. The yield of the final step was 21 mg, 75%.

MS [M-CF₃COO]⁺: 460

¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m, 8H), 1.80-2.10 (m, 6H), 2.28 (m, 1H), 2.75-2.85 (m, 3H), 3.10-3.55 (m, 7H), 3.80-3.90 (m, 1H), 5.05-5.15 (m, 1H), 6.20 (s, 1H, OH),
25 6.90-6.95 (m, 1H), 6.95-7.05 (m, 2H), 7.10-7.20 (m, 1H), 7.35-7.45 (m, 2H).

Example 24

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide

The title compound was synthesised according to methods c, and a from the Intermediate I-15a. The yield of the final step was 338 mg, 92%.

MS [M-Br]⁺: 456; mp: 75°C

35 ¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.75-2.10 (m, 4H), 2.27-2.35 (m, 1H), 2.70-2.90 (m, 1H), 3.30-3.68 (m, 5H), 3.68-3.83 (m, 2H), 3.92-4.10 (m, 1H), 4.32-4.50 (m,

2H), 5.10-5.20 (m, 1H), 6.20 (s, 1H, OH), 6.90-7.05 (m, 4H), 7.10-7.20 (m, 1H), 7.30-7.42 (m, 3H).

5 **Example 25**

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 23. The yield of the final step was 17 mg, 64%.

10 MS [M-CF₃COO]⁺: 440

¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m, 8H), 1.80-2.10 (m, 4H), 2.32 (m, 1H), 2.75-2.85 (m, 1H), 2.95-3.05 (m, 2H), 3.20-3.50 (m, 6H), 3.50-3.65 (m, 1H), 3.85-3.95 (m, 1H), 5.10-5.20 (m, 1H), 6.22 (s, 1H, OH), 6.95-7.05 (m, 1H), 7.10-7.20 (m, 1H), 7.20-7.40 (m, 5H), 7.40-7.55 (m, 1H)

15

Example 26

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

20 The title compound was synthesised as in Example 23. The yield of the final step was 20 mg, 74%.

MS [M-CF₃COO]⁺: 454

¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m, 8H), 1.75-2.05 (m, 6H), 2.28 (m, 1H), 2.55-2.65 (m, 2H), 2.75-2.85 (m, 1H), 3.10-3.40 (m, 6H), 3.40-3.55 (m, 1H), 3.77-3.87 (m, 1H), 25 5.05-5.15 (m, 1H), 6.20 (s, 1H, OH), 6.95-7.0 (m, 1H), 7.10-7.15 (m, 1H), 7.20-7.35 (m, 5H), 7.38-7.42 (m, 1H).

Example 27

30 **(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenylallyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised as in Example 23. The yield of the final step was 3 mg, 12%.

MS [M-CF₃COO]⁺: 452

35 ¹H-NMR (DMSO-d₆): δ 1.25-1.60 (m, 8H), 1.80-2.10 (m, 4H), 2.31 (m, 1H), 2.72-2.85 (m, 1H), 3.12-3.22 (m, 1H), 3.22-3.45 (m, 3H), 3.45-3.60 (m, 1H), 3.82-3.92 (m, 1H),

3.95-4.10 (m, 2H), 5.10-5.20 (m, 1H), 6.20 (s, 1H, OH), 6.35-6.50 (m, 1H), 6.82-6.95 (m, 2H), 7.10-7.15 (m, 1H), 7.25-7.47 (m, 4H), 7.55-7.62 (m, 2H)

5 **Example 28**

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-fluorophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 23. The yield of the final step was 15 mg, 52%.

10 MS [M-CF₃COO]⁺: 488

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.80-2.05 (m, 4H), 2.05-2.17 (m, 2H), 2.30 (m, 1H), 2.75-2.90 (m, 1H), 3.17-3.27 (m, 1H), 3.27-3.60 (m, 6H), 3.82-3.95 (m, 1H), 3.97-4.05 (m, 2H), 5.14 (m, 1H), 6.22 (s, 1H, OH), 6.92-7.05 (m, 3H), 7.10-7.20 (m, 3H), 7.40-7.55 (m, 1H).

15

Example 29

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(4-oxo-4-thien-2-ylbutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

20 The title compound was synthesised as in Example 23. The yield of the final step was 4 mg, 14%.

MS [M-CF₃COO]⁺: 488

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.70-2.05 (m, 6H), 2.30 (m, 1H), 2.75-2.90 (m, 1H), 3.05-3.12 (m, 2H), 3.15-3.60 (m, 7H), 3.80-3.92 (m, 1H), 5.13 (m, 1H), 6.22 (s, 1H, OH), 6.98-7.02 (m, 1H), 7.12-7.18 (m, 1H), 7.25-7.30 (m, 1H), 7.40-7.55 (m, 1H), 7.95-8.0 (m, 1H), 8.02-8.07 (m, 1H).

25

Example 30

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[4-(4-fluorophenyl)-4-oxobutyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

30

The title compound was synthesised as in Example 23. The yield of the final step was 9 mg, 29%.

MS [M-CF₃COO]⁺: 500

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.75-2.05 (m, 6H), 2.30 (m, 1H), 2.75-2.90 (m, 1H), 3.05-3.60 (m, 9H), 3.80-3.95 (m, 1H), 5.14 (m, 1H), 6.22 (s, 1H, OH), 6.98-7.02 (m, 1H), 7.12-7.20 (m, 1H), 7.35-7.45 (m, 3H), 8.02-8.12 (m, 2H).

35

Example 31

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-(3-hydroxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 23. The yield of the final step was

5 14 mg, 48%.

MS [M-CF₃COO]⁺: 486

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.80-2.0 (m, 4H), 2.0-2.20 (m, 2H), 2.30 (m, 1H), 2.75-2.90 (m, 1H), 3.17-3.25 (m, 1H), 3.25-3.60 (m, 6H), 3.82-3.92 (m, 1H), 3.94-4.02 (m, 2H), 5.14 (m, 1H), 6.21 (s, 1H, OH), 6.30-6.42 (m, 3H), 6.95-7.10 (m, 2H),
10 7.12-7.20 (m, 1H), 7.20-7.45 (m, 1H), 9.47 (s, 1H, OH).

Example 32

1-(2-Benzoyloxyethyl)-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

15 The title compound was synthesised as in Example 23. The yield of the final step was 10 mg, 35%.

MS [M-CF₃COO]⁺: 470

¹H-NMR (DMSO-d₆): δ 1.25-1.62 (m, 8H), 1.80-2.07 (m, 4H), 2.30 (m, 1H), 2.75-2.85 (m, 1H), 3.0-3.65 (m, 7H), 3.75-4.0 (m, 3H), 4.50 (s, 2H), 5.10-5.17 (m, 1H), 6.21 (s, 1H, OH), 6.95-7.0 (m, 1H), 7.10-7.17 (m, 1H), 7.27-7.45 (m, 6H).
20

Example 33

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-1-(3-o-tolyloxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

25 The title compound was synthesised as in Example 23. The yield of the final step was 13 mg, 45%.

MS [M-CF₃COO]⁺: 484

¹H-NMR (DMSO-d₆): δ 1.23-1.65 (m, 8H), 1.80-2.05 (m, 4H), 2.05-2.20 (m, 5H), 2.31 (m, 1H), 2.75-2.90 (m, 1H), 3.15-3.25 (m, 1H), 3.27-3.60 (m, 6H), 3.85-3.95 (m, 1H), 3.97-4.05 (m, 2H), 5.15 (m, 1H), 6.22 (s, 1H, OH), 6.83-6.93 (m, 2H), 6.98-7.02 (m, 1H),
30 7.12-7.20 (m, 3H), 7.40-7.46 (m, 1H).

Example 34

1-[3-(3-Cyanophenoxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
35

The title compound was synthesised as in Example 23. The yield of the final step was 11 mg, 32%.

MS [M-CF₃COO]⁺: 495

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.75-2.05 (m, 4H), 2.05-2.20 (m, 2H), 2.30 (m, 1H), 2.75-2.90 (m, 1H), 3.20-3.25 (m, 1H), 3.25-3.60 (m, 6H), 3.82-3.95 (m, 1H), 4.05-4.15 (m, 2H), 5.07-5.20 (m, 1H), 6.20 (s, 1H, OH), 6.95-7.05 (m, 1H), 7.12-7.20 (m, 1H), 7.25-7.35 (m, 1H), 7.40-7.57 (m, 4H)

Example 35

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 23. The yield of the final step was 10 mg, 26%.

MS [M-CF₃COO]⁺: 520, (described in experimental section, method b).

¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m, 8H), 1.80-2.10 (m, 4H), 2.20-2.37 (m, 3H), 2.75-2.92 (m, 1H), 3.20-3.65 (m, 7H), 3.90-4.05 (m, 1H), 4.15-4.30 (m, 2H), 5.15-5.22 (m, 1H), 6.24 (s, 1H, OH), 6.95-7.05 (m, 2H), 7.15-7.20 (m, 1H), 7.40-7.60 (m, 5H), 7.85-7.95 (m, 1H), 8.20-8.25 (m, 1H)

Example 36

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(methylephenylamino)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 23. The yield of the final step was 12 mg, 35%.

MS [M-CF₃COO]⁺: 483

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.75-2.02 (m, 6H), 2.28 (m, 1H), 2.75-2.85 (m, 1H), 2.87 (s, 3H), 3.09-3.14 (m, 1H), 3.15-3.55 (m, 8H), 3.75-3.87 (m, 1H), 5.05-5.15 (m, 1H), 6.20 (s, 1H, OH), 6.60-6.70 (m, 1H), 6.70-6.77 (m, 2H), 6.92-7.0 (m, 1H), 7.10-7.25 (m, 3H), 7.35-7.45 (m, 1H).

Example 37

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 23. The yield of the final step was 7 mg, 22%.

MS [M-CF₃COO]⁺: 486

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.75-2.02 (m, 6H), 2.27 (m, 1H), 2.75-2.90 (m, 1H), 2.95-3.05 (m, 2H), 3.07-3.15 (m, 1H), 3.15-3.52 (m, 1H), 3.75-3.87 (m, 1H), 5.05-5.15 (m, 1H), 6.20 (s, 1H, OH), 6.95-7.0 (m, 1H), 7.12-7.17 (m, 1H), 7.20-7.30 (m, 1H), 7.30-7.45 (m, 5H).

5

Example 38

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-1-(4-oxo-4-phenylbutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

10 The title compound was synthesised as in Example 23. The yield of the final step was 9 mg, 26%.

MS [M-CF₃COO]⁺: 482

¹H-NMR (DMSO-d₆): δ 1.30-1.67 (m, 8H), 1.72-2.10 (m, 6H), 2.30 (m, 1H), 2.75-2.90 (m, 1H), 3.10-3.60 (m, 9H), 3.85-3.95 (m, 1H), 5.10-5.20 (m, 1H), 6.23 (s, 1H, OH),
15 6.95-7.05 (m, 1H), 7.12-7.20 (m, 1H), 7.40-7.47 (m, 1H), 7.52-7.60 (m, 2H), 7.62-7.72 (m, 1H), 7.95-8.05 (m, 1H).

Example 39

20 **(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-1-[3-(2,4,6-trimethylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised as in Example 23. The yield of the final step was 13 mg, 35%.

MS [M-CF₃COO]⁺: 512

25 ¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m, 8H), 1.80-2.02 (m, 4H), 2.02-2.25 (m, 11H), 2.32 (m, 1H), 2.75-2.90 (m, 1H), 3.23-3.28 (m, 1H), 3.28-3.62 (m, 6H), 3.65-3.80 (m, 2H), 3.85-3.97 (m, 1H), 5.10-5.20 (m, 1H), 6.22 (s, 1H, OH), 6.82 (s, 2H), 6.97-7.05 (m, 1H), 7.12-7.20 (m, 1H), 7.40-7.47 (m, 1H)

30

Example 40

1-[3-(2-Chlorophenoxy)-propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

35 The title compound was synthesised as in Example 23. The yield of the final step was 13 mg, 36%.

MS [M-CF₃COO]⁺: 505

¹H-NMR (DMSO-d₆): 1.30-1.65 (m, 8H), 1.75-2.05 (m, 4H), 2.05-2.25 (m, 2H), 2.31 (m, 1H), 2.75-2.90 (m, 1H), 3.19-3.23 (m, 1H), 3.23-3.62 (m, 1H), 3.85-4.0 (m, 1H), 4.07-4.15 (m, 2H), 5.10-5.20 (m, 1H), 6.22 (s, 1H, OH), 6.92-7.05 (m, 2H), 7.12-7.22 (m, 2H), 7.27-7.37 (m, 1H), 7.40-7.50 (m, 2H).

5

Example 41

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(3-trifluoromethylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

10 The title compound was synthesised as in Example 23. The yield of the final step was 13 mg, 33%.

MS [M-CF₃COO]⁺: 538

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.75-2.05 (m, 4H), 2.05-2.25 (m, 2H), 2.31 (m, 1H), 2.75-2.90 (m, 1H), 3.20-3.25 (m, 1H), 3.25-3.62 (m, 6H), 3.82-3.97 (m, 1H),
15 4.05-4.20 (m, 2H), 5.10-5.20 (m, 1H), 6.22 (s, 1H, OH), 6.95-7.05 (m, 1H), 7.12-7.20 (m, 1H), 7.22-7.30 (m, 2H), 7.30-7.37 (m, 1H), 7.40-7.47 (m, 1H), 7.50-7.62 (m, 1H).

Example 42

20 **1-[3-(Biphenyl-4-yloxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised as in Example 23. The yield of the final step was 11 mg, 30%.

MS [M-CF₃COO]⁺: 546

25 ¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.77-2.05 (m, 4H), 2.05-2.25 (m, 2H), 2.31 (m, 1H), 2.75-2.92 (m, 1H), 3.20-3.23 (m, 1H), 3.23-3.62 (m, 6H), 3.85-3.97 (m, 1H), 4.05-4.15 (m, 2H), 5.10-5.20 (m, 1H), 6.22 (s, 1H, OH), 6.95-7.10 (m, 3H), 7.12-7.20 (m, 1H), 7.27-7.37 (m, 1H), 7.40-7.50 (m, 3H), 7.55-7.70 (m, 4H).

30

Example 43

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(2,4-difluorophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 23. The yield of the final step was
35 10 mg, 28%.

MS [M-CF₃COO]⁺: 506

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.75-2.05 (m, 4H), 2.05-2.15 (m, 2H), 2.30 (m, 1H), 2.82 (m, 1H), 3.17-3.28 (m, 1H), 3.28-3.47 (m, 5H), 3.60 (m, 1H), 3.82-3.95 (m, 1H), 4.05-4.15 (m, 2H), 5.14 (m, 1H), 6.22 (s, 1H, OH), 6.95-7.10 (m, 2H), 7.12-7.38 (m, 3H), 7.40-7.45 (m, 1H).

5

Example 44

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-methoxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 23. The yield of the final step was

10 11 mg, 32%.

MS [M-CF₃COO]⁺: 500

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.75-2.15 (m, 6H), 2.30 (m, 1H), 2.82 (m, 1H), 3.18-3.25 (m, 1H), 3.25-3.45 (m, 5H), 3.45-3.60 (m, 1H), 3.70 (s, 3H), 3.82-3.92 (m, 1H), 3.92-4.02 (m, 2H), 5.14 (m, 1H), 6.22 (s, 1H, OH), 6.88 (m, 4H), 6.98-7.02 (m, 1H), 7.15-7.16 (m, 1H), 7.42-7.44 (m, 1H).

15

Example 45

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(5,6,7,8-tetrahydronaphthalen-2-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane

20 trifluoroacetate

The title compound was synthesised as in Example 23. The yield of the final step was 14 mg, 38%.

MS [M-CF₃COO]⁺: 524

¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m, 8H), 1.65-1.75 (m, 4H), 1.75-2.20 (m, 6H), 2.30 (m, 1H), 2.50-2.70 (m, 4H), 2.75-2.95 (m, 1H), 3.17-3.25 (m, 1H), 3.25-3.45 (m, 5H), 3.45-3.60 (m, 1H), 3.80-3.92 (m, 1H), 3.92-4.02 (m, 2H), 5.14 (m, 1H), 6.22 (s, 1H, OH), 6.60-6.70 (m, 2H), 6.95-7.02 (m, 2H), 7.15-7.20 (m, 1H), 7.42-7.45 (m, 1H).

25

Example 46

30 **1-[3-(Benzo[1,3]dioxol-5-yloxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised as in Example 23. The yield of the final step was 13 mg, 38%.

MS [M-CF₃COO]⁺: 514

35 ¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.75-2.15 (m, 6H), 2.30 (m, 1H), 2.75-2.90 (m, 1H), 3.15-3.25 (m, 1H), 3.25-3.42 (m, 5H), 3.42-3.60 (m, 1H), 3.82-3.92 (m, 1H),

3.92-4.0 (m, 2H), 5.12 (m, 1H), 5.97 (s, 2H); 6.20 (s, 1H, OH), 6.36-6.40 (m, 1H), 6.64-6.65 (m, 1H), 6.81-6.84 (m, 1H), 6.98-7.02 (m, 1H), 7.15-7.17 (m, 1H), 7.42-7.44 (m, 1H).

5

Example 47

1-[3-(2-Carbamoyl-phenoxy)-propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-yl-acetoxy]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 23. The yield of the final step was

10 13 mg, 36%.

MS [M-CF₃COO]⁺: 513

¹H-NMR (DMSO-d₆): δ 1.27-1.65 (m, 8H), 1.80-2.07 (m, 4H), 2.12-2.27 (m, 2H), 2.31 (m, 1H), 2.82 (m, 1H), 3.17-3.25 (m, 1H), 3.25-3.45 (m, 5H), 3.45-3.60 (m, 1H), 3.82-3.92 (m, 1H), 4.10-4.17 (m, 2H), 5.15 (m, 1H), 6.23 (s, 1H, OH), 6.98-7.16 (m, 4H),

15 7.42-7.50 (m, 2H), 7.50-7.55 (bs, 2H), 7.68-7.72 (m, 1H).

Example 48

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(3-dimethylaminophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

20 The title compound was synthesised as in Example 23. The yield of the final step was 14 mg, 40%.

MS [M-CF₃COO]⁺: 513

25 ¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.70-2.20 (m, 6H), 2.30 (m, 1H), 2.75-2.95 (m, 7H), 3.15-3.65 (m, 7H), 3.80-4.05 (m, 3H), 5.14 (m, 1H), 6.15-6.30 (m, 3H), 6.32-6.36 (m, 1H), 6.95-7.22 (m, 3H), 7.40-7.45 (m, 1H).**Example 49**

1-[3-(4-Acetylaminophenoxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

30 The title compound was synthesised as in Example 23. The yield of the final step was 12 mg, 34%.

MS [M-CF₃COO]⁺: 527

35 ¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.75-2.20 (m, 9H), 2.30 (m, 1H), 2.82 (m, 1H), 3.17-3.28 (m, 1H), 3.28-3.45 (m, 5H), 3.45-3.60 (m, 1H), 3.84-3.92 (m, 1H), 3.96-4.02 (m, 2H), 5.13 (m, 1H), 6.22 (s, 1H, OH), 6.86-6.90 (m, 2H), 6.98-7.02 (m, 1H), 7.15-7.17 (m, 1H), 7.42-7.44 (m, 1H), 7.48-7.52 (m, 2H), 9.85 (s, 1H, NH(CO)).

Example 50**(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-****5 methoxycarbonylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised as in Example 23. The yield of the final step was 14 mg, 37%.

MS [M-CF₃COO]⁺: 528

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.80-2.05 (m, 4H), 2.05-2.25 (m, 2H), 2.31
10 (m, 1H), 2.82 (m, 1H), 3.17-3.28 (m, 1H), 3.28-3.45 (m, 5H), 3.45-3.60 (m, 1H), 3.82 (s, 3H), 3.82-3.95 (m, 1H), 4.10-4.15 (m, 2H), 5.14 (m, 1H), 6.22 (s, 1H, OH), 6.95-7.10 (m, 3H), 7.15-7.17 (m, 1H), 7.43-7.45 (m, 1H), 7.92-7.97 (m, 2H).

Example 51

15 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-nitrophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 23. The yield of the final step was 15 mg, 41%.

MS [M-CF₃COO]⁺: 515

¹H-NMR (DMSO-d₆): δ 1.27-1.65 (m, 8H), 1.77-2.07 (m, 4H), 2.10-2.27 (m, 2H), 2.31
20 (m, 1H), 2.82 (m, 1H), 3.17-3.28 (m, 1H), 3.28-3.45 (m, 5H), 3.45-3.60 (m, 1H), 3.85-3.95 (m, 1H), 4.15-4.25 (m, 2H), 5.15 (m, 1H), 6.22 (s, 1H, OH), 6.97-7.02 (m, 1H), 7.14-7.18 (m, 3H), 7.42-7.45 (m, 1H), 8.22-8.27 (m, 2H).

25 Example 52**(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-****hydroxymethylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised as in Example 23. The yield of the final step was 13 mg, 36%.

30 MS [M-CF₃COO]⁺: 500

¹H-NMR (DMSO-d₆): δ 1.25-1.65 (m, 8H), 1.77-2.05 (m, 4H), 2.05-2.20 (m, 2H), 2.30
(m, 1H), 2.82 (m, 1H), 3.17-3.60 (m, 7H), 3.82-3.95 (m, 1H), 3.95-4.05 (m, 2H), 4.35-4.45 (m, 2H), 5.05-5.11 (t, 1H, OH), 5.11-5.20 (m, 1H), 6.22 (s, 1H, OH), 6.86-6.95 (m, 2H), 6.95-7.05 (m, 1H), 7.15-7.17 (m, 1H), 7.22-7.26 (m, 2H), 7.42-7.44 (m, 1H).

35

Example 53

(3R)-3-[(2R)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-phenoxypropyl]-1-azoniabicyclo[2.2.2]octane bromide

The title compound was synthesised according to methods d, and a from the Intermediate I-15b. The yield of the final step was 1.2 g (73%).

mp: 181°C. MS : [M-Br]⁺: 470.

¹H-NMR (DMSO-d₆): δ 1.30-1.70 (m, 8H), 1.70-1.80 (m, 2H), 1.80-2.05 (m, 2H), 2.05-2.30 (m, 3H), 2.80-2.95 (m, 1H), 3.25-3.62 (m, 7H), 3.87-4.0 (m, 1H), 4.0-4.10 (m, 2H), 5.10-5.20 (m, 1H), 6.20 (s, 1H, OH), 6.95-7.05 (m, 4H), 7.15-7.25 (m, 1H), 7.25-7.37 (m, 2H), 7.42-7.45 (m, 1H).

Example 54

(3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods d, and b from the Intermediate I-15b. The yield of the final step was 15 mg (54%).

MS [M-CF₃COO]⁺: 460

¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m 8H), 1.65-1.80 (m, 2H), 1.80-2.10 (m, 4H), 2.21 (m, 1H), 2.77-2.90 (m, 3H), 3.15-3.40 (m, 6H), 3.40-3.55 (m, 1H), 3.80-3.92 (m, 1H), 5.05-5.18 (m, 1H), 6.20 (s, 1H, OH), 6.92-6.96 (m, 1H), 6.96-7.02 (m, 2H), 7.12-7.20 (m, 1H), 7.36-7.40 (m, 1H), 7.40-7.46 (m, 1H)

Example 55

(3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 54. The yield of the final step was 16 mg, 58%.

MS [M-CF₃COO]⁺: 456.

¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m, 8H), 1.75-1.85 (m, 2H), 1.85-2.05 (m, 2H), 2.23 (m, 1H), 2.75-2.90 (m, 1H), 3.40-3.57 (m, 4H), 3.57-3.70 (m, 1H), 3.70-3.80 (m, 2H), 3.97-4.10 (m, 1H), 4.37-4.47 (m, 2H), 5.10-5.18 (m, 1H), 6.20 (s, 1H, OH), 6.92-7.05 (m, 4H), 7.10-7.18 (m, 1H), 7.30-7.38 (m, 2H), 7.38-7.44 (m, 1H),

Example 56

(3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 54. The yield of the final step was 13 mg, 50%.

MS [M-CF₃COO]⁺: 440.

¹H-NMR (DMSO-d₆): δ 1.35-1.65 (m, 8H), 1.65-1.85 (m, 2H), 1.85-2.05 (m, 2H), 2.25 (m, 1H), 2.85-2.92 (m, 1H), 2.95-3.10 (m, 2H), 3.30-3.50 (m, 6H), 3.50-3.65 (m, 1H), 3.85-4.0 (m, 1H), 5.12-5.20 (m, 1H), 6.21 (s, 1H, OH), 6.95-7.05 (m, 1H), 7.15-7.20 (m, 1H), 7.25-7.40 (m, 5H), 7.40-7.47 (m, 1H).

Example 57

(3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 54. The yield of the final step was 14 mg, 53%.

MS [M-CF₃COO]⁺: 454.

¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m, 8H), 1.65-1.77 (m, 2H), 1.77-2.05 (m, 4H), 2.21 (s, 1H), 2.55-2.65 (m, 2H), 2.75-2.92 (m, 1H), 3.15-3.40 (m, 6H), 3.40-3.55 (m, 1H), 3.80-3.90 (m, 1H), 5.06-5.16 (m, 1H), 6.19 (s, 1H, OH), 6.95-7.02 (m, 1H), 7.12-7.18 (m, 1H), 7.20-7.36 (m, 5H), 7.38-7.46 (m, 1H).

Example 58

(3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-(3-phenylallyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 54. The yield of the final step was 7 mg, 26%.

MS [M-CF₃COO]⁺: 452.

¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m, 8H), 1.75-2.05 (m, 4H), 2.24 (m, 1H), 2.75-2.90 (m, 1H), 3.25-3.45 (m, 4H), 3.45-3.55 (m, 1H), 3.72-3.95 (m, 1H), 4.0-4.15 (m, 2H), 5.10-5.17 (m, 1H), 6.19 (s, 1H, OH), 6.40-6.55 (m, 1H), 6.82-6.70 (m, 2H), 7.12-7.17 (m, 1H), 7.30-7.45 (m, 4H), 7.55-7.62 (m, 2H)

Example 59

(3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl-1-[3-(4-fluorophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

5 The title compound was synthesised as in Example 54. The yield of the final step was 19 mg, 64%.

MS [M-CF₃COO]⁺: 488.

¹H-NMR (DMSO-d₆): δ 1.30-1.65 (m, 8H), 1.65-1.80 (m, 2H), 1.80-2.05 (m, 2H), 2.05-2.20 (m, 2H), 2.23 (m, 1H), 2.80-2.95 (m, 1H), 3.20-3.60 (m, 7H), 3.85-3.95 (m, 1H),
10 3.97-4.07 (m, 2H), 5.14 (m, 1H), 6.20 (s, 1H, OH), 6.90-7.05 (m, 3H), 7.10-7.20 (m, 3H), 7.40-7.47 (m, 1H).

Example 60

(3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxyl-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

15 The title compound was synthesised according to methods d, and b from Intermediate I-14a. The yield of the final step was 4 mg, 15%.

MS [M-CF₃COO]⁺: 454

¹H-NMR (DMSO-d₆): δ 1.45-1.67 (m, 9H), 1.67-1.80 (m, 1H), 1.80-2.05 (m, 2H), 2.05-2.22 (m, 3H), 2.85-2.95 (m, 1H), 3.20-3.55 (m, 7H), 3.85-3.95 (m, 1H), 4.0-4.10 (m, 2H), 5.10-5.20 (m, 1H), 6.03 (s, 1H), 6.40-6.45 (m, 2H), 6.90-7.0 (m, 3H), 7.27-7.35 (m, 2H), 7.62 (m, 1H)

Example 61

(3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxyl-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 60. The yield of the final step was 2 mg, 7%.

30 MS [M-CF₃COO]⁺: 444

¹H-NMR (DMSO-d₆): δ 1.40-1.66 (m, 9H), 1.66-2.10 (m, 5H), 2.17 (m, 1H), 2.78-2.90 (m, 3H), 3.14-3.50 (m, 7H), 3.80-3.90 (m, 1H), 5.10-5.18 (m, 1H), 6.02 (s, 1H), 6.38-6.46 (m, 2H), 6.92-7.02 (m, 2H), 7.36-7.40 (m, 1H), 7.60 (m, 1H).

35

Example 62

(3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-phenylethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 60. The yield of the final step was 4 mg, 17%.

MS [M-CF₃COO]⁺: 424

¹H-NMR (DMSO-d₆): δ 1.40-2.05 (m, 12H), 2.20 (m, 1H), 2.88 (m, 1H), 2.95-3.05 (m, 2H), 3.20-3.60 (m, 7H), 3.85-3.95 (m, 1H), 5.18 (m, 1H), 6.03 (s, 1H, OH), 6.40-6.45 (m, 2H), 7.25-7.40 (m, 5H), 7.62 (m, 1H).

Example 63

(3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 60. The yield of the final step was 10 mg, 36.2%.

MS [M-CF₃COO]⁺: 438

¹H-NMR (DMSO-d₆): δ 1.35-2.05 (m, 14H), 2.16 (m, 1H), 2.55-2.65 (m, 2H), 2.75-2.95 (m, 1H), 3.10-3.55 (m, 7H), 3.77-3.92 (m, 1H), 5.05-5.15 (m, 1H), 6.02 (s, 1H, OH), 6.35-6.45 (m, 2H), 7.17-7.40 (m, 5H), 7.60 (m, 1H).

Example 64

(3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods d, and b from Intermediate I-14b. The yield of the final step was 12 mg, 46%.

MS [M-CF₃COO]⁺: 454

¹H-NMR (DMSO-d₆): δ 1.40-1.60 (m, 8H), 1.75-2.05 (m, 4H), 2.05-2.20 (m, 2H), 2.30 (m, 1H), 2.75-2.87 (m, 1H), 3.10-3.60 (m, 7H), 3.85-3.95 (m, 1H), 3.97-4.07 (m, 2H), 5.10-5.17 (m, 1H), 6.04 (s, 1H), 6.40-6.50 (m, 2H), 6.90-7.0 (m, 3H), 7.27-7.37 (m, 2H), 7.60-7.65 (m, 1H).

Example 65

(3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 64. The yield of the final step was 14 mg, 55%.

MS [M-CF₃COO]⁺: 444

¹H-NMR (DMSO-d₆): δ 1.40-1.65 (m, 8H), 1.75-2.10 (m, 6H), 2.27 (m, 1H), 2.70-2.90 (m, 3H), 3.0-3.55 (m, 7H), 3.77-3.82 (m, 1H), 5.05-5.15 (m, 1H), 6.03 (s, 1H), 6.40-6.45 (m, 2H), 6.90-7.05 (m, 2H), 7.35-7.42 (m, 1H), 7.55-7.65 (m, 1H).

Example 66

(3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 64. The yield of the final step was 15 mg, 57%.

MS [M-CF₃COO]⁺: 424

¹H-NMR (DMSO-d₆): δ 1.30-2.40 (m, 13H), 2.75-2.85 (m, 1H), 2.95-3.05 (m, 2H), 3.10-3.75 (m, 7H), 3.85-4.0 (m, 1H), 5.05-5.15 (m, 1H), 6.02 (s, 1H, OH), 6.44 (m, 2H), 7.20-7.40 (m, 5H), 7.63 (m, 1H).

Example 67

(3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 64. The yield of the final step was 10 mg, 38%.

MS [M-CF₃COO]⁺: 438

¹H-NMR (DMSO-d₆): δ 1.40-1.60 (m, 8H), 1.75-2.05 (m, 6H), 2.27 (m, 1H), 2.55-2.60 (m, 2H), 2.79 (m, 1H), 3.04-3.10 (m, 1H), 3.12-3.40 (m, 5H), 3.40-3.52 (m, 1H), 3.80-3.90 (m, 1H), 5.10 (m, 1H), 6.02 (s, 1H, OH), 6.40 (m, 2H), 7.20-7.35 (m, 5H), 7.58 (m, 1H)

Example 68

(3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods d, and b from the Intermediate I-16a. The yield of the final step was 28 mg, 100%.

MS [M-CF₃COO]⁺: 478.

¹H-NMR (DMSO-d₆): δ 0.95-1.18 (m, 4H), 1.18-1.65 (m, 5H), 1.70-2.05 (m, 5H), 2.05-2.17 (m, 2H), 2.17-2.30 (m, 2H), 3.15-3.25 (m, 1H), 3.25-3.5 (m, 6H), 3.75-3.90 (m, 1H), 3.95-4.07 (m, 2H), 5.05-5.15 (m, 1H), 5.78 (s, 1H, OH), 6.90-7.0 (m, 3H), 7.25-7.45 (m, 5H), 7.55-7.65 (m, 2H).

5

Example 69

(3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxyl-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

10 The title compound was synthesised as in Example 68. The yield of the final step was 22 mg, 78%.

MS [M-CF₃COO]⁺: 468.

¹H-NMR (DMSO-d₆): δ 0.95-1.18 (m, 4H), 1.18-1.65 (m, 5H), 1.70-2.0 (m, 7H), 2.20 (m, 2H), 2.75-2.85 (m, 2H), 3.05-3.15 (m, 1H), 3.15-3.50 (m, 6H), 3.70-3.85 (m, 1H),
15 5.05-5.15 (m, 1H), 5.76 (s, 1H, OH), 6.90-7.05 (m, 2H), 7.20-7.45 (m, 4H), 7.55-7.65 (m, 2H)

Example 70

20 **(3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxyl-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised as in Example 68. The yield of the final step was 14 mg, 50%.

MS [M-CF₃COO]⁺: 448.

25 ¹H-NMR (DMSO-d₆): δ 0.97-1.15 (m, 3H), 1.15-2.05 (m, 11H), 2.15-2.30 (m, 2H), 2.90-3.05 (m, 2H), 3.20-3.30 (m, 1H), 3.30-3.50 (m, 5H), 3.50-3.62 (m, 1H), 3.82-3.92 (m, 1H), 5.15 (m, 1H), 5.78 (s, 1H, OH), 7.25-7.45 (m, 8H), 7.58-7.64 (m, 2H).

Example 71

30 **(3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxyl-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods d, and b from the Intermediate I-16b. The yield of the final step was 18 mg, 63%.

35 MS [M-CF₃COO]⁺: 478.

¹H-NMR (DMSO-d₆): δ 0.95-1.17 (m, 4H), 1.20-2.0 (m, 10H), 2.02-2.35 (m, 4H), 3.15-3.55 (m, 7H), 3.80-3.90 (m, 1H), 3.97-4.10 (m, 2H), 5.05-5.12 (m, 1H), 5.75 (s, 1H, OH), 6.90-7.02 (m, 3H), 7.25-7.45 (m, 5H), 7.57-7.67 (m, 2H)

5

Example 72

(3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 71. The yield of the final step was 19 mg, 66 %.

MS [M-CF₃COO]⁺: 468.

¹H-NMR (DMSO-d₆): δ 0.95-1.17 (m, 4H), 1.20-2.10 (m, 12H), 2.15-2.35 (m, 2H), 2.75-2.97 (m, 2H), 3.10-3.37 (m, 6H), 3.37-3.55 (m, 1H), 3.75-3.87 (m, 1H), 5.05-5.12 (m, 1H), 5.74 (s, 1H, OH), 6.90-7.05 (m, 2H), 7.22-7.45 (m, 4H), 7.55-7.67 (m, 2H).

15

Example 73

(3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 71. The yield of the final step was 16 mg, 58 %.

MS [M-CF₃COO]⁺: 448.

¹H-NMR (DMSO-d₆): δ 0.98-1.15 (m, 3H), 1.20-2.05 (m, 11H), 2.20-2.35 (m, 2H), 2.90-3.10 (m, 2H), 3.20-3.50 (m, 6H), 3.50-3.60 (m, 1H), 3.80-3.92 (m, 1H), 5.12 (m, 1H), 5.75 (s, 1H, OH), 7.25-7.40 (m, 8H), 7.60-7.65 (m, 1H).

25

Example 74

(3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods c, and b from the intermediate I-4a. The yield of the final step was 3.8 mg, 6.2%.

MS [M-CF₃COO]⁺: 468.

¹H-NMR (DMSO-d₆): δ 1.0-1.50 (m, 7H), 1.50-2.05 (m, 7H), 2.05-2.35 (m, 4H), 3.15-3.65 (m, 7H), 3.82-3.95 (m, 1H), 4.0-4.1 (m, 2H), 5.16 (m, 1H), 5.99 (s, 1H, OH), 6.40-6.45 (m, 2H), 6.90-7.0 (m, 3H), 7.25-7.35 (m, 2H), 7.64 (m, 1H).

35

Example 75

(3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phen-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 74. The yield of the final step was 3.6 mg , 6 %.

MS [M-CF₃COO]⁺: 458.

¹H-NMR (DMSO-d₆): δ 1.0-1.45 (m, 7H), 1.50-2.10 (m, 9H), 2.15-2.30 (m, 2H), 2.75-2.90 (m, 2H), 3.10-3.55 (m, 7H), 3.77-3.92 (m, 1H), 5.13 (m, 1H), 5.98 (s, 1H, OH), 6.36-6.46 (m, 2H), 6.92-7.02 (m, 2H), 7.36-7.40 (m, 1H), 7.62 (m, 1H).

Example 76

(3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 74. The yield of the final step was 2.4 mg , 4.2 %.

MS [M-CF₃COO]⁺: 438.

¹H-NMR (DMSO-d₆): δ 1.0-1.45 (m, 7H), 1.50-2.05 (m, 7H), 2.20-2.35 (m, 2H), 2.92-3.15 (m, 2H), 3.20-3.65 (m, 7H), 3.85-3.95 (m, 1H), 5.18 (m, 1H), 6.0 (s, 1H, OH), 6.37-6.47 (m, 2H), 7.25-7.45 (m, 5H), 7.64 (m, 1H).

Example 77

(3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 74. The yield of the final step was 2.8 mg , 4.8 %.

MS [M-CF₃COO]⁺: 452.

¹H-NMR (DMSO-d₆): δ 0.95-1.50 (m, 7H), 1.50-2.10 (m, 9H), 2.15-2.32 (m, 2H), 2.55-2.65 (m, 2H), 3.10-3.60 (m, 7H), 3.77-3.80 (m, 1H), 5.12 (m, 1H), 5.98 (s, 1H, OH), 6.36-6.46 (m, 2H), 7.18-7.40 (m, 5H), 7.62 (m, 1H).

Example 78

(3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods c, and b from the Intermediate I-4b. The yield of the final step was 3.0 mg , 5%.

MS [M-CF₃COO]⁺: 468.

¹H-NMR (DMSO-d₆): 1.0-1.45 (m, 7H), 1.55-1.75 (m, 3H), 1.80-2.05 (m, 4H), 2.05-2.25 (m, 3H), 2.30 (m, 1H), 3.10-3.20 (m, 1H), 3.20-3.60 (m, 1H), 3.85-3.95 (m, 1H), 3.95-4.10 (m, 2H), 5.16 (m, 1H), 5.99 (s, 1H, OH), 6.40-6.50 (m, 2H), 6.90-7.0 (m, 3H), 7.25-7.38 (m, 2H), 7.64 (m, 1H).

5

Example 79

(3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxyl]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

10 The title compound was synthesised as in Example 78. The yield of the final step was 9.1 mg , 33.1 %.

MS [M-CF₃COO]⁺: 458.

¹H-NMR (DMSO-d₆): δ 0.95-1.55 (m, 7H), 1.55-2.10 (m, 9H), 2.10-2.40 (m, 2H), 2.75-2.95 (m, 2H), 3.0-3.12 (m, 1H), 3.12-3.70 (m, 6H), 3.80-3.95 (m, 1H), 5.14 (m, 1H), 6.0 (s, 1H, OH), 6.35-6.55 (m, 2H), 6.90-7.10 (m, 2H), 7.35-7.45 (m, 1H), 7.60-7.70 (m, 1H).

15

Example 80

(3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxyl]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

20

The title compound was synthesised as in Example 78. The yield of the final step was 3.6 mg , 6 %.

MS [M-CF₃COO]⁺: 438.

¹H-NMR (DMSO-d₆): δ 1.0-1.45 (m, 7H), 1.55-1.80 (m, 3H), 1.80-2.10 (m, 4H), 2.12-2.28 (m, 1H), 2.30 (m, 1H), 2.90-3.05 (m, 2H), 3.15-3.25 (m, 1H), 3.25-3.50 (m, 5H), 3.50-3.65 (m, 1H), 3.85-3.95 (m, 1H), 5.18 (m, 1H), 6.0 (s, 1H, OH), 6.38-6.48 (m, 2H), 7.24-7.40 (m , 5H), 7.65 (m, 1H).

25

Example 81

(3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxyl]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

30

The title compound was synthesised as in Example 78. The yield of the final step was 5.8 mg , 10 %.

35 MS [M-CF₃COO]⁺: 452.

¹H-NMR (DMSO-d₆): δ 1.0-1.42 (m, 7H), 1.55-1.77 (m, 3H), 1.77-2.05 (m, 6H), 2.18 (m, 1H), 2.27 (m, 1H), 2.55-2.65 (m, 2H), 3.02-3.12 (m, 1H), 3.12-3.30 (m, 6H), 3.77-3.90 (m, 1H), 5.13 (m, 1H), 5.98 (s, 1H, OH), 6.40 (m, 2H), 7.20-7.35 (m, 5H), 7.61 (m, 1H).

5 **Example 82**

(3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods c, and b from Intermediate I-5a. The yield of the final step was 9.4 mg, 15.6 %.

10 MS [M-CF₃COO]⁺: 464

¹H-NMR (DMSO-d₆): δ 1.10-1.70 (m, 8H), 1.70-2.02 (m, 4H), 2.05-2.15 (m, 2H), 2.24 (m, 1H), 2.90-2.97 (m, 1H), 3.15-3.25 (m, 1H), 3.25-3.60 (m, 6H), 3.75-3.92 (m, 1H), 3.95-4.10 (m, 2H), 5.10 (m, 1H), 5.86 (s, 1H, OH), 6.90-7.0 (m, 3H), 7.20-7.40 (m, 5H), 7.56-7.66 (m, 2H).

15

Example 83

(3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 82. The yield of the final step was 20 5.0 mg, 8.8 %.

MS [M-CF₃COO]⁺: 434

¹H-NMR (DMSO-d₆): δ 1.12-1.70 (m, 8H), 1.75-2.05 (m, 4H), 2.26 (m, 1H), 2.87-3.05 (m, 3H), 3.15-3.62 (m, 7H), 3.80-3.92 (m, 1H), 5.13 (m, 1H), 5.86 (s, 1H, OH), 7.24-7.44 (m, 8H), 7.56-7.66 (m, 2H).

25

Example 84

(3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 82. The yield of the final step was 30 3.2 mg, 5.1%.

MS [M-CF₃COO]⁺: 480

¹H-NMR (DMSO-d₆): δ 1.12-1.65 (m, 8H), 1.65-2.0 (m, 6H), 2.21 (m, 1H), 2.85-3.15 (m, 4H), 3.15-3.55 (m, 6H), 3.70-3.85 (m, 1H), 5.06 (m, 1H), 5.83 (s, 1H, OH), 7.20-7.46 (m, 8H), 7.54-7.64 (m, 2H).

35

Example 85

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods c, and b from Intermediate I-5b. The yield of the final step was 7.8 mg, 12.9 %.

MS [M-CF₃COO]⁺: 464

¹H-NMR (DMSO-d₆): δ 1.15-1.35 (m, 2H), 1.35-2.0 (m, 10H), 2.0-2.30 (m, 3H), 2.95-3.10 (m, 1H), 3.20-3.60 (m, 7H), 3.80-3.95 (m, 1H), 3.97-4.10 (m, 2H), 5.09 (m, 1H), 5.84 (s, 1H, OH), 6.90-7.0 (m, 3H), 7.24-7.44 (m, 5H), 7.60-7.70 (m, 2H).

Example 86

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 85. The yield of the final step was 5.2 mg, 9.2%.

MS [M-CF₃COO]⁺: 434

¹H-NMR (DMSO-d₆): δ 1.12-1.32 (m, 2H), 1.32-2.05 (m, 10H), 2.20 (m, 1H), 2.90-3.10 (m, 3H), 3.20-3.62 (m, 7H), 3.82-3.97 (m, 1H), 5.12 (m, 1H), 5.85 (s, 1H, OH), 7.22-7.45 (m, 8H), 7.60-7.70 (m, 2H).

Example 87

(3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 85. The yield of the final step was 4.0 mg, 6.4%.

MS [M-CF₃COO]⁺: 480

¹H-NMR (DMSO-d₆): δ 1.12-1.32 (m, 2H), 1.32-1.70 (m, 8H), 1.70-2.0 (m, 4H), 2.16 (m, 1H), 2.92-3.05 (m, 3H), 3.15-3.60 (m, 7H), 3.75-3.87 (m, 1H), 5.04 (m, 1H), 5.82 (s, 1H, OH), 7.20-7.44 (m, 8H), 7.58-7.68 (m, 2H).

Example 88

(3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods c, and b from intermediate I-6a. The yield of the final step was 3.2 mg, 5.1%.

MS [M-CF₃COO]⁺: 454

¹H-NMR (DMSO-d₆): δ 1.0-1.45 (m, 7H), 1.55-1.80 (m, 3H), 1.80-2.20 (m, 7H), 2.25-2.35 (m, 1H), 3.20-3.28 (m, 1H), 3.28-3.42 (m, 5H), 3.42-3.55 (m, 1H), 3.85-3.95 (m, 1H), 4.01-4.05 (m, 2H), 5.17 (m, 1H), 6.16 (s, 1H, OH), 6.92-7.03 (m, 4H), 7.13-7.15 (m, 1H), 7.28-7.34 (m, 2H), 7.42-7.45 (m, 1H).

Example 89

(3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 88. The yield of the final step was 3.4 mg, 5.8%.

MS [M-CF₃COO]⁺: 454

¹H-NMR (DMSO-d₆): δ 1.0-1.45 (m, 7H), 1.55-1.80 (m, 3H), 1.80-2.15 (m, 5H), 2.32 (m, 1H), 2.95-3.05 (m, 2H), 3.20-3.52 (m, 6H), 3.52-3.68 (m, 1H), 3.85-3.95 (m, 1H), 5.20 (m, 1H), 6.16 (s, 1H, OH), 7.0-7.04 (m, 1H), 7.10-7.15 (m, 1H), 7.25-7.40 (m, 5H), 7.43-7.46 (m, 1H).

Example 90

1-[3-(3-Cyanophenoxy)propyl]-(3R)-3-[(2S)-2-cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 88. The yield of the final step was 8.4 mg, 12.7%.

MS [M-CF₃COO]⁺: 509

¹H-NMR (DMSO-d₆): δ 1.0-1.45 (m, 7H), 1.55-1.70 (m, 3H), 1.70-2.20 (m, 7H), 2.31 (m, 1H), 3.20-3.27 (m, 1H), 3.25-3.50 (m, 5H), 3.45-3.60 (m, 1H), 3.85-3.95 (m, 1H), 4.05-4.15 (m, 2H), 5.18 (m, 1H), 6.16 (s, 1H, OH), 7.0-7.03 (m, 1H), 7.13-7.15 (m, 1H), 7.28-7.31 (m, 1H), 7.43-7.46 (m, 3H), 7.50-7.55 (m, 1H).

Example 91

(3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods c, and b from Intermediate I-6b (The Intermediate I-6b used herein contains approx. 25% of I-6a, see previous description in method c). The yield of the final step was 3.0 mg, 4.8%.

MS [M-CF₃COO]⁺: 484

¹H-NMR (DMSO-d₆): δ 1.0-1.50 (m, 7H), 1.55-2.22 (m, 10H), 2.25-2.35 (m, 1H), 3.20-3.45 (m, 5H), 3.45-3.60 (m, 1H), 3.85-3.95 (m, 1H), 4.0-4.1 (m, 2H), 5.17 (m, 1H), 6.14 (and 6.16) (s, 1H, OH, mixture of diastereomers aprox. 75:25), 6.93-7.03 (m, 4H), 7.13-7.17 (m, 1H), 7.28-7.35 (m, 2H), 7.42-7.45 (m, 1H).

5

Example 92

(3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 91. The yield of the final step was 2.6 mg, 4.4%.

10

MS [M-CF₃COO]⁺: 454

¹H-NMR (DMSO-d₆): δ 1.50-1.70 (m, 7H), 1.55-2.20 (m, 8H), 2.28 (m, 1H), 2.95-3.10 (m, 2H), 3.20-3.52 (m, 6H), 3.52-3.65 (m, 1H), 3.85-3.97 (m, 1H), 5.15-5.25 (m, 1H), 6.14 (and 6.16), (s, 1H, OH, mixture of diastereomers aprox. 75:25), 6.98-7.04 (m, 1H), 7.13-7.16 (m, 1H), 7.25-7.40 (m, 5H), 7.43-7.46 (m, 1H).

15

Example 93

1-[3-(3-Cyanophenoxy)propyl]-(3R)-3-[(2R)-2-cyclohexyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 91. The yield of the final step was 5.0 mg, 7.5 %.

20

MS [M-CF₃COO]⁺: 509

¹H-NMR (DMSO-d₆): δ 1.0-1.50 (m, 7H), 1.55-2.05 (m, 7H), 2.05-2.22 (m, 3H), 2.22-2.35 (m, 1H), 3.20-3.50 (m, 6H), 3.45-3.60 (m, 1H), 3.80-3.95 (m, 1H), 4.10-4.15 (m, 2H), 5.17 (m, 1H), 6.14 (and 6.16) (s, 1H, OH, mixture of diastereomers aprox. 75:25), 7.0-7.03 (m, 1H), 7.13-7.16 (m, 1H), 7.28-7.32 (m, 1H), 7.43-7.46 (m, 3H), 7.50-7.55 (m, 1H).

25

Example 94

(3R)-3-(2-Hydroxy-4-phenyl-2-thien-2-ylbutanoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as a mixture of diastereomers according to methods d, and b from Intermediate I-21. The yield of the final step was 4.8 mg, 15.8 %.

35

MS [M-CF₃COO]⁺: 506

¹H-NMR (DMSO-d₆): δ 1.75-2.05 (m, 3H), 2.05-2.20 (m, 2H), 2.20-2.35 (m, 2H), 2.37-2.70 (m, 4H), 3.20-3.65 (m, 7H), 3.82-3.95 (m, 1H), 4.0-4.1 (m, 1H), 5.12 (m, 1H), 6.58 (s, 1H, OH), 6.90-7.0 (m, 3H), 7.0-7.08 (m, 1H), 7.14-7.24 (m, 4H), 7.24-7.36 (m, 4H), 7.46-7.52 (m, 1H).

5

Example 95

(3S)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide

The title compound was synthesised according to methods c, and a from the Intermediate I-7. The yield of the final step was 250 mg, 87.1%.

MS [M-Br]⁺: 470

¹H-NMR (DMSO-d₆): δ (Same description as in Example 53)

Example 96

15 **(3S)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods c, and b from the Intermediate I-7. The yield of the final step was 11.1 mg, 40.2%.

MS [M-CF₃COO]⁺: 460

20 ¹H-NMR (DMSO-d₆): δ (Same description as in Example 54)

Example 97

(3S)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

25 The title compound was synthesised as in Example 96. The yield of the final step was 11.3 mg, 41.4%.

MS [M-CF₃COO]⁺: 454.

¹H-NMR (DMSO-d₆): δ (Same description as in Example 57).

Example 98

4-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxyl]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods c, and b from the Intermediate I-8. The yield was 9.4 mg, 34.6%.

35 MS [M-CF₃COO]⁺: 468

¹H-NMR (DMSO-d₆): δ 0.90-1.65 (m, 9H), 1.70-1.80 (m, 1H), 1.90-2.05 (m, 2H), 2.05-2.20 (m, 1H), 2.18-2.35 (m, 6H), 2.75-2.90 (m, 2H), 3.10-3.25 (m, 2H), 3.45-3.70 (m,

6H), 5.60 (s, 1H, OH), 6.90-6.92 (m, 1H), 6.95-7.02 (m, 1H), 7.20-7.45 (m, 4H), 7.50-7.60 (m, 2H).

5 Example 99

4-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised as in Example 98. The yield of the final step was 8.1 mg, 29.2%.

10 MS [M-CF₃COO]⁺: 478

¹H-NMR (DMSO-d₆): δ 0.90-1.65 (m, 9H), 1.70-1.80 (m, 1H), 2.05-2.20 (m, 3H), 2.20-2.40 (m, 6H), 3.20-3.45 (m, 2H), 3.50-3.75 (m, 6H), 3.95-4.10 (m, 2H), 5.60 (s, 1H, OH), 6.85-7.05 (m, 3H), 7.20-7.45 (m, 5H), 7.50-7.65 (m, 2H).

15

((*): Configuration not assigned)

20 The following examples illustrate pharmaceutical compositions according to the present invention and procedures for their preparation.

Example 100

Preparation of a pharmaceutical composition: tablets

Formulation:

25	Compound of the present invention	5.0 mg
	Lactose	113.6 mg
	Microcrystalline cellulose	28.4 mg
	Light silicic anhydride	1.5 mg
	Magnesium stearate	1.5 mg

30

Using a mixer machine, 15 g of the compound of the present invention was mixed with 340.8 g of lactose and 85.2 g of microcrystalline cellulose. The mixture was subjected to compression moulding using a roller compactor to give a flake-like compressed material. The flake-like compressed material was pulverized using a hammer mill, and 35 the pulverized material was screened through a 20 mesh screen. A 4.5 g portion of light silicic anhydride and 4.5 g of magnesium stearate were added to the screened material and mixed. The mixer product was subjected to a tablets making machine

equipped with a die/punch system of 7.5 mm in diameter, thereby obtaining 3,000 tablets each having 150 mg in weight.

Example 101

5 Preparation of a pharmaceutical composition: tablets coated

Formulation:

	Compound of the present invention	5.0 mg
	Lactose	95.2 mg
	Corn starch	40.8 mg
10	Polyvinylpyrrolidone K25	7.5 mg
	Magnesium stearate	1.5 mg
	Hydroxypropylcellulose	2.3 mg
	Polyethylene glycol 6000	0.4 mg
	Titanium dioxide	1.1 mg
15	Purified talc	0.7 mg

Using a fluidized bed granulating machine, 15 g of the compound of the present invention was mixed with 285.6 g of lactose and 122.4 g of corn starch. Separately, 22.5 g of polyvinylpyrrolidone was dissolved in 127.5 g of water to prepare a binding solution. Using a fluidized bed granulating machine, the binding solution was sprayed on the above mixture to give granulates. A 4.5 g portion of magnesium stearate was added to the obtained granulates and mixed. The obtained mixture was subjected to a tablet making machine equipped with a die/punch biconcave system of 6.5 mm in diameter, thereby obtaining 3,000 tablets, each having 150 mg in weight.

25 Separately, a coating solution was prepared by suspending 6.9 g of hydroxypropylmethylcellulose 2910, 1.2 g of polyethylene glycol 6000, 3.3 g of titanium dioxide and 2.1 g of purified talc in 72.6 g of water. Using a High Coated, the 3,000 tablets prepared above were coated with the coating solution to give film-coated tablets, each having 154.5 mg in weight.

30

Example 102

Preparation of a pharmaceutical composition: liquid inhalant

Formulation:

	Compound of the present invention	400 µg
35	Physiological saline	1 ml

A 40 mg portion of the compound of the present invention was dissolved in 90 ml of physiological saline, and the solution was adjusted to a total volume of 100 ml with the same saline solution, dispensed in 1 ml portions into 1 ml capacity ampoule and then sterilized at 115° for 30 minutes to give liquid inhalant.

5

Example 103**Preparation of a pharmaceutical composition: powder inhalant**

Formulation:

Compound of the present invention	200 µg
10 Lactose	4,000 µg

A 20 g portion of the compound of the present invention was uniformly mixed with 400 g of lactose, and a 200 mg portion of the mixture was packed in a powder inhaler for exclusive use to produce a powder inhalant.

15

Example 104**Preparation of a pharmaceutical composition: inhalation aerosol.**

Formulation:


Compound of the present invention	200 µg
20 Dehydrated (Absolute) ethyl alcohol USP	8,400 µg
1,1,1,2-Tetrafluoroethane (HFC-134A)	46,810 µg

The active ingredient concentrate is prepared by dissolving 0.0480 g of the compound of the present invention in 2.0160 g of ethyl alcohol. The concentrate is added to an appropriate filling apparatus. The active ingredient concentrate is dispensed into aerosol container, the headspace of the container is purged with Nitrogen or HFC-134A vapor (purging ingredients should not contain more than 1 ppm oxygen) and is sealed with valve. 11.2344 g of HFC-134A propellant is then pressure filled into the sealed container.

25

CLAIMS

1. A compound of formula (I):



(I)

5

wherein B is a phenyl ring, a 5 to 10 membered heteroaromatic group containing one or more heteroatoms, or a naphthalenyl, 5,6,7,8-tetrahydronaphthalenyl, benzo[1,3]dioxolyl, or biphenyl group;

R^1 , R^2 and R^3 each independently represent a hydrogen or halogen atom, or a hydroxy group, a phenyl group, $-OR^7$, $-SR^7$, $-NR^7R^8$, $-NHCOR^7$, $-CONR^7R^8$, $-CN$, $-NO_2$, $-COOR^7$ or $-CF_3$ group, or a straight or branched, substituted or unsubstituted lower alkyl group, wherein R^7 and R^8 each independently represent a hydrogen atom, a straight or branched lower alkyl group, or together form an alicyclic ring; or R^1 and R^2 together form an aromatic or alicyclic ring or a heterocyclic group.

15

n is an integer from 0 to 4;

A represents a group selected from $-CH_2-$, $-CH=CR^9-$, $-CR^9=CH-$, $-CR^9R^{10}-$, $-CO-$, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$ and NR^9 , wherein R^9 and R^{10} each independently represent a hydrogen atom, a straight or branched lower alkyl group, or together form an alicyclic ring;


20

m is an integer from 0 to 8, provided that when $m=0$, A is not $-CH_2-$;

p is an integer from 1 to 2 and the substitution in the azonia bicyclic ring may be in the 2, 3 or 4 position including all possible configurations of the asymmetric carbons;

25


R^4 represents a group of structure:



30

wherein R^{11} represents a hydrogen or halogen atom, a peroxy group, an alkoxy group, a nitro group, a cyano group, $-\text{CO}_2R^{12}$ or $-\text{NR}^{12}R^{13}$, wherein R^{12} and R^{13} are identical or different and are selected from hydrogen and straight or branched lower alkyl groups, or a straight or branched, substituted or unsubstituted lower alkyl group;

R^5 represents an alkyl group of 1 to 7 carbon atoms, an alkenyl group containing 2 to 7 carbon atoms, an alkynyl group containing 2 to 7 carbon atoms, a cycloalkyl group of 3 to 7 carbon atoms, or a group of formula



wherein $q = 1$ or 2 and R^{11} is as defined above;

R^6 represents a hydrogen atom, a hydroxy group, a methyl group or a $-\text{CH}_2\text{OH}$ group; and

X^- represents a pharmaceutically acceptable anion of a mono or polyvalent acid.

2. A compound according to claim 1, wherein p is 2.
3. A compound according to claim 1 or claim 2, wherein R^4 represents an unsubstituted phenyl, 2-thienyl, 3-thienyl, 2-furyl or 3-furyl group.
4. A compound according to any one of the preceding claims, wherein R^5 represents a cyclopentyl, cyclohexyl, pentyl, allyl, vinyl, propynyl, benzyl or phenethyl group.
5. A compound according to any one of the preceding claims, wherein the group $-\text{O}-\text{CO}-\text{C}(R^4)(R^5)(R^6)$ represents a group selected from 2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl; 2,3-diphenylpropionyloxy; 2-hydroxymethyl-2,3-diphenylpropionyloxy; 2-hydroxy-2,3-diphenylpropionyloxy; 2-hydroxy-3-phenyl-2-thien-2-ylpropionyloxy; 2-hydroxy-2-thien-2-ylpent-4-enoyloxy; 2-hydroxy-2-thien-2-ylheptanoyloxy; 2-hydroxy-2-thien-2-ylpent-3-ynoyloxy; 2-hydroxy-2-thien-2-ylbut-3-enoyloxy; 2-cyclopentyl-2-fur-2-yl-2-hydroxyacetoxyl; 2-cyclohexyl-2-hydroxy-2-phenylacetoxyl; 2-cyclohexyl-2-fur-2-yl-2-hydroxyacetoxyl; 2-

cyclopentyl-2-hydroxy-2-phenylacetoxy, 2-cyclohexyl-2-hydroxy-2-thien-2-ylacetoxy and 2-hydroxy-4-phenyl-2-thien-2-ylbutanoyloxy

- 5 6. A compound according to any one of the preceding claims, wherein B represents a phenyl, pyrrolyl, thienyl, furyl, biphenyl, naphthalenyl, 5,6,7,8-tetrahydronaphthalenyl, benzo[1,3]dioxolyl, imidazolyl or benzothiazolyl group.
- 10 7. A compound according to claim 6, wherein B represents a phenyl, pyrrolyl or thienyl group.
8. A compound according to any one of the preceding claims wherein R¹, R² and R³ each independently represent a hydrogen or halogen atom or a hydroxy, methyl,

15 tert-butyl, -CH₂OH, 3-hydroxypropyl, -OMe, -NMe₂, -NHCOMe, -CONH₂, -CN, -NO₂, -COOMe or -CF₃ group.
9. A compound according to claim 8 wherein R¹, R² and R³ each independently represent a hydrogen or halogen atom or a hydroxy group.
- 20 10. A compound according to any one of the preceding claims, wherein n= 0 or 1; m is an integer from 1 to 6; and A represents a -CH₂-, -CH=CH-, -CO-, -NMe-, -O- or -S- group.
- 25 11. A compound according to claim 10, wherein m is 1, 2 or 3 and A represents a -CH₂-, -CH=CH-, -O- or -S- group.
12. A compound according to any one of the preceding claims, wherein the azoniabicyclo group is substituted on the nitrogen atom with a group selected

30 from 3-phenoxypropyl, 2-phenoxyethyl, 3-phenylallyl, phenethyl, 3-phenylpropyl, 3-(3-hydroxyphenoxy)propyl, 3-(4-fluorophenoxy)propyl, 3-thien-2-ylpropyl, 4-oxo-4-thien-2-ylbutyl, 2-benzyloxyethyl, 3-o-tolyloxypropyl, 3-(3-cyanophenoxy)propyl, 3-(methylphenylamino)propyl, 3-phenylsulfanylpropyl, 4-oxo-4-phenylbutyl, 3-(2-chlorophenoxy)propyl, 3-(2,4-difluorophenoxy)propyl, 3-

35 (4-methoxyphenoxy)propyl, 3-(benzo[1,3]dioxol-5-yloxy)propyl.

13. A compound according to any one of the preceding claims, wherein X⁻ represents a chloride, bromide, trifluoroacetate or methanesulphonate anion.
14. A compound according to any one of the preceding claims, wherein the azoniabicyclic group is substituted in the 3-position.
15. A compound according to claim 14, wherein the substituent at the 3-position of the azoniabicyclic group has R configuration.
16. A compound according to claim 14, wherein the substituent at the 3-position of the azoniabicyclic group has S configuration.
17. A compound according to any one of the preceding claims, wherein the carbon substituted by R⁴, R⁵ and R⁶ has R configuration.
18. A compound according to any one of claims 1 to 16 wherein the carbon substituted by R⁴, R⁵ and R⁶ has S configuration.
19. A compound according to any one of the preceding claims, which is a single isomer.
20. A compound according to claim 1 which is:

(3R)-3-(2,3-Diphenylpropionyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

(3R)-3-(2,3-Diphenylpropionyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

(3R)-3-(2-Hydroxymethyl-2,3-diphenylpropionyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

(3R)-3-(2-Hydroxymethyl-2,3-diphenylpropionyloxy)-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

(3R)-3-(2-Hydroxy-2,3-diphenylpropionyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide

(3R)-3-(2-Hydroxy-2,3-diphenylpropionyloxy)-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide

(3R)-3-(2-Hydroxy-3-phenyl-2-thien-2-ylpropionyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

- (3R)-3-(2-Hydroxy-2-phenyl-2-thien-2-ylpropionyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-2-thien-2-ylpent-4-enoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- 5 (3R)-3-(2-Hydroxy-2-thien-2-ylpent-4-enoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-2-thien-2-ylpent-4-enoyloxy)-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-(2-Hydroxy-2-thien-2-ylheptanoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 10 (3R)-3-(2-Hydroxy-2-thien-2-ylheptanoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-2-thien-2-ylpent-3-ynoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 (3R)-3-(2-Hydroxy-2-thien-2-ylpent-3-ynoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-2-thien-2-ylbut-3-enoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-2-thien-2-ylbut-3-enoyloxy)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenylallyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-fluorophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 35 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(4-oxo-4-thien-2-ylbutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate


- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[4-(4-fluorophenyl)-4-oxobutyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(3-hydroxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 1-(2-Benzyl-oxyethyl)-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-o-tolyloxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 10 1-[3-(3-Cyanophenoxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(methylphenylamino)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(4-oxo-4-phenylbutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(2,4,6-trimethylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 1-[3-(2-Chlorophenoxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(3-trifluoromethylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 1-[3-(Biphenyl-4-yloxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(2,4-difluorophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-methoxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(5,6,7,8-tetrahydronaphthalen-2-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 1-[3-(Benzo[1,3]dioxol-5-yloxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 35 1-[3-(2-Carbamoylphenoxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(3-dimethylaminophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 1-[3-(4-Acetylaminophenoxy)propyl]-(3R)-3-[(2S)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-methoxycarbonylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-nitrophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-hydroxymethylphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 10 (3R)-3-[(2R)-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-(3-phenylallyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxyl]-1-[3-(4-fluorophenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 (3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxyl]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxyl]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxyl]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 (3R)-3-[(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxyl]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxyl]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 35 (3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxyl]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate


- (3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-phenethyl-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenylpropyl)-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-thien-2-ylpropyl)-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-phenethyl-1-
10 azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-thien-2-ylpropyl)-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 (3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-phenethyl-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenoxypropyl)-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-thien-2-ylpropyl)-1-
20 azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-phenethyl-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenylpropyl)-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 (3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenoxypropyl)-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-thien-2-ylpropyl)-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-phenethyl-1-
30 azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetoxy]-1-(3-phenylpropyl)-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-
azoniabicyclo[2.2.2]octane trifluoroacetate
- 35 (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-phenethyl-1-
azoniabicyclo[2.2.2]octane trifluoroacetate

- (3R)-3-[(2R)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 10 (3R)-3-[(2S)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 1-[3-(3-Cyanophenoxy)propyl]-(3R)-3-[(2S)-2-cyclohexyl-2-hydroxy-2-thien-2-ylacetoxy]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 (3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(2R)-2-Cyclohexyl-2-hydroxy-2-thien-2-ylacetoxy]-1-phenethyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 1-[3-(3-Cyanophenoxy)propyl]-(3R)-3-[(2R)-2-cyclohexyl-2-hydroxy-2-thien-2-ylacetoxy]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 (3R)-3-(2-Hydroxy-4-phenyl-2-thien-2-ylbutanoyloxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- 25 (3S)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-3-[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetoxy]-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 4-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 4-[(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate


21. A process for producing a compound of formula (I):



wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , p , m , n , A , B and X are as defined in any one of claims 1 to 18, which process comprises quaternising the nitrogen atom of the azoniabicyclic ring of a compound of formula (III):



wherein R^4 , R^5 , R^6 and p are as defined above, with an alkylating agent of formula (II):



wherein R^1 , R^2 , R^3 , m , n , A and B are as defined above, and W represents a leaving group.


15

22. A process according to claim 21, wherein W represents a group X as defined in claims 1 or 13.

20

23. A process according to claim 21 or claim 22, wherein the resulting reaction mixture is purified by solid phase extraction.

24. A compound of formula (III)



(III)

wherein p, R₄, R₅ and R₆ are as defined in claim 1, the substituent on the azabicyclo group is at position 3 or 4 and when it is at position 3 this substituted carbon has an enantiomerically pure R or S configuration,

5

provided that when R₄ is a 3-thienyl group and R₅ is a cyclohexyl group R₆ is not a hydroxy group.

25. A compound according to claim 24, wherein the substituent on the azabicyclo group is at position 3 with R configuration.

10

26. A compound according to claim 24, wherein the substituent on the azabicyclo group is at position 3 with S configuration.

15 27. A compound according to any one of claims 24 to 26 wherein the carbon substituted by R⁴, R⁵ and R⁶ has R configuration.

28. A compound according to any one of claims 24 to 26 wherein the carbon substituted by R⁴, R⁵ and R⁶ has S configuration.

20

29. A compound according to any one of claims 24 to 28, which is a single isomer.

30. A compound according to claim 24

25

which is:

(2R)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester;

(2S)-2-Cyclopentyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester;

30

(2R)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester;

(2S)-2-Cyclohexyl-2-fur-2-yl-2-hydroxyacetic acid (3R)-1-azabicyclo[2.2.2]oct-3-yl ester;
(2R)-2-Cyclohexyl-2-hydroxy-2-phenylacetic acid 1-azabicyclo[2.2.2]oct-4-yl ester

5

31. A pharmaceutical composition comprising a compound according to any one of claims 1 to 20 in admixture with a pharmaceutically acceptable carrier or diluent.
- 10 32. A compound according to any one of claims 1 to 20, or a pharmaceutical composition according to claim 31, for therapeutic use in a method of treatment of the human or animal body by therapy.
- 15 33. Use of a compound according to any one of claims 1 to 20, or a pharmaceutical composition according to claim 31, in the manufacture of a medicament for the treatment of a respiratory, urological or gastrointestinal disease or disorder.
- 20 34. A method of treating a respiratory, urological or gastrointestinal disease or disorder, which method comprises administering to a human or animal patient in need of treatment an effective non-toxic amount of a compound according to any one of claims 1 to 20 or a pharmaceutical composition according to claim 31.
- 25 35. A combination product comprising
(i) a compound according to any one of claims 1 to 20; and
(iii) another compound effective in the treatment of a respiratory, urological or gastrointestinal disease or disorder
for simultaneous, separate or sequential use.
- 30 36. A combination product according to claim 35 comprising
(i) a compound according to any one of claims 1 to 20; and
(iv) a β_2 agonist, steroid, antiallergic drug, phosphodiesterase IV inhibitor and/or leukotriene D4 (LTD4) antagonist
for simultaneous, separate or sequential use in the treatment of a respiratory
35 disease.

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
11 July 2002 (11.07.2002)

PCT

(10) International Publication Number
WO 02/053564 A3(51) International Patent Classification⁷: C07D 453/02,
A61K 31/439, A61P 43/00(ES). BUIL ALBERO, Maria, Antonia [ES/ES]; Paris,
50, 1^o, 4a, E-08029 Barcelona (ES).

(21) International Application Number: PCT/EP01/15168

(74) Agents: SRINIVASAN, Ravi, Chandran et al.; J.A.
Kemp & Co., 14 South Square, Gray's Inn, London WC1R
5JJ (GB).(22) International Filing Date:
20 December 2001 (20.12.2001)

(25) Filing Language: English

(26) Publication Language: English




(30) Priority Data:
P0003130 28 December 2000 (28.12.2000) ES(71) Applicant (for all designated States except US): ALMI-
RALL PRODESFARMA AG [CH/CH]; Lindenhof, Dorf-
strasse 38, CH-6341 Baar (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PRAT QUINONES,
Maria [ES/ES]; Calle Andrea Doria, 2, 1^o 1a, E-08003
Barcelona (ES). FERNANDEZ FORNER, Maria, Do-
lors [ES/ES]; Roger de Flor 221, 5^o, 4a, E-08025 Barcelona(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

[Continued on next page]

(54) Title: QUINUCLIDINE DERIVATIVES AND THEIR USE AS M3 ANTAGONISTS



(57) Abstract: A compound of formula (I), wherein B is a phenyl ring, a 5 to 10 membered heteroaromatic group containing one or more heteroatoms, or a naphthalenyl, 5, 6, 7, 8- tetrahydronaphthalenyl, benzo[1,3]dioxolyl, or biphenyl group; A represents a group selected from -CH₂-, -CH=CR⁹-, -CR⁹=CH-, -CR⁹R¹⁰-, -CO-, -O-, -S-, -S(O)-, -S(O)₂- and NR⁹, R⁴ represents a group of structure: (II) R⁵ represents an-alkyl group of 1 to 7 carbon atoms, an alkenyl group containing 2 to 7 carbon atoms, an alkyl group containing 2 to 7 carbon atoms, a cycloalkyl group of 3 to 7 carbon atoms, or a group of formula (a) wherein q = 1 or 2 and R¹¹ is as defined above; R⁶ represents a hydrogen atom, a hydroxy group, a methyl group or a -CH₂OH group; and X⁻ represents a pharmaceutically acceptable anion of a mono or polyvalent acid.



Published:

— *with international search report*

(88) Date of publication of the international search report:

31 October 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/15168

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D453/02 A61K 31/439 A61P43/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 863 141 A (BANYU PHARMACEUTICAL CO., LTD.) 9 September 1998 (1998-09-09) page 21, line 29 - line 33; claims	1, 31-33
A	EP 0 930 298 A (BANYU PHARMACEUTICAL CO., LTD.) 21 July 1999 (1999-07-21) page 50, line 15 - line 24; claims	1, 31-33
P, A	WO 01 04118 A (ALMIRALL PRODESPHARMA S.A.) 18 January 2001 (2001-01-18) cited in the application claims	1, 31-33

☐ Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

2 July 2002

Date of mailing of the international search report

12/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 01/15168

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 34 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/15168

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 863141	A	09-09-1998	AT 205490 T	15-09-2001
			AU 7145996 A	30-04-1997
			DE 69615214 D1	18-10-2001
			DE 69615214 T2	27-06-2002
			EP 0863141 A1	09-09-1998
			US 6130232 A	10-10-2000
			CA 2234619 A1	17-04-1997
			WO 9713766 A1	17-04-1997
EP 930298	A	21-07-1999	AU 716050 B2	17-02-2000
			AU 3635197 A	25-02-1998
			BG 103114 A	30-11-1999
			BR 97111108 A	17-08-1999
			EE 9900038 A	16-08-1999
			EP 0930298 A1	21-07-1999
			JP 3063164 B2	12-07-2000
			NO 990472 A	01-02-1999
			NZ 333842 A	25-05-2001
			SK 12299 A3	16-05-2000
			CN 1226888 A	25-08-1999
			CZ 9900331 A3	14-07-1999
			HR 970426 A1	31-08-1998
			HU 9902381 A2	29-11-1999
			WO 9805641 A1	12-02-1998
			JP 3282617 B2	20-05-2002
			JP 2000178231 A	27-06-2000
			JP 3282618 B2	20-05-2002
			JP 2000169449 A	20-06-2000
			PL 331431 A1	19-07-1999
			TR 9900204 T2	21-01-2000
			TR 200001482 T2	21-11-2000
			US 6040449 A	21-03-2000
			US 5948792 A	07-09-1999
			ZA 9706813 A	11-02-1998
WO 0104118	A	18-01-2001	ES 2165768 A1	16-03-2002
			AU 6433000 A	30-01-2001
			BR 0012434 A	02-04-2002
			WO 0104118 A2	18-01-2001
			EP 1200431 A2	02-05-2002
			NO 20020180 A	13-03-2002

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.